

***Hypoxia-inducible factor 3A* gene expression and methylation in adipose tissue is related to adipose tissue dysfunction**

Dissertation

zur Erlangung des akademischen Grades Dr. med.

an der Medizinischen Fakultät

der Universität Leipzig

eingereicht von:

Susanne Erna Margarete Pfeiffer

Geburtsdatum / Geburtsort:

02.04.1989 in München

Angefertigt am 25.08.2016 in Leipzig

Betreuer:

Prof. Dr. med. Matthias Blüher

Beschluss über die Verleihung des Doktorgrades vom: 21.03.2017

Table of contents

List of abbreviations	4
1 Introduction	7
1.1 Obesity and its associated pathomechanisms	7
1.1.1 Obesity: a global health burden	7
1.1.2 Adipose tissue	8
1.1.3 Development of hypoxia in adipose tissue	9
1.1.4 Inflammation in adipose tissue	10
1.2 The transcription factor family of “hypoxia inducible factor”	11
1.2.1 The function of hypoxia inducible factors	11
1.2.2 The role of hypoxia inducible factor 3 α	12
1.3 Genetics and epigenetics.....	14
1.3.1 Epigenetic mechanisms.....	14
1.3.2 The methylation of DNA	15
1.3.3 Epigenetics in association with obesity	15
1.3.4 Single nucleotide polymorphisms in obesity	16
1.4 Main hypothesis	16
1.5 Aim of this study.....	17
2 Manuscript of the publication	18
3 Summary.....	28
4 Supplemental material.....	32
4.1 Material and methods.....	32
4.2 Results.....	34
5 Declaration of the independent formulation of the work	37
6 Curriculum vitae	38
7 Acknowledgement.....	40
8 References.....	41

List of abbreviations

ANT1 / ANT2	adenine nucleotide translocase 1 / 2
AT	adipose tissue
% body fat	percentage of body fat
°C	degree celcius
µg	microgram
µkat	microkatal
µl	microliter
µmol	micromol
adi	adipocyte
adj.	adjusted
ALAT	alanine aminotransferase
ASAT	aspartate aminotransferase
AU	arbitrary unit
bHLH	basal Helix-loop-helix domain
BMI	body mass index
CBP	calcium response element binding protein
cDNA	complementary DNA
c/EBPα	CCAAT/enhancer binding protein α
cm	centimetre
CpG	cytosine-phosphate-guanine dinucleotide
CRP	C-reactive protein
CT	computed tomography
CT ratio	visceral fat area / subcutaneous fat area
C-TAD	C-terminal transactivation domain
DNA	deoxyribonucleic acid
dl	deciliter
DNMT	DNA methyltransferase
dNTP	deoxyribose nucleoside triphosphate
ENCODE	encyclopedia of DNA elements
EWAS	epigenome-wide association studies
FAM	6-Carboxy-Fluorescein
FD	fat distribution
FFA	free fatty acid
FIH	factor inhibiting HIF
FPG	fasting plasma glucose
FPI	fasting plasma insulin
fT3	free triiodthyronine

ft4	free tetraiodothyronine
g	gram
GEO	gene expression omnibus
gGT	gamma-glutamyl transferase
GIR	glucose infusion rate
HbA1c	glycohemoglobin
HDL-C	high density lipoprotein cholesterol
HFD	high fat diet
HIF1α / HIF2α / HIF3α / HIFβ	Hypoxia-Inducible Factor 1 α / 2 α / 3 α / β
HIF-p4H	HIF prolyl 4-hydroxylase
HPRT1	hypoxanthine guanine phosphoribosyltransferase 1
HRE	hypoxia responsive element
IL-1b / IL-6b	interleukin 1b / 6b
kg	kilogram
l	liter
LDL-C	low density lipoprotein cholesterol
ln	natural logarithm
LXXLL	peptide recognition motif 'Leu-Xaa-Xaa-Leu-Leu'
LZIP	leucine zipper
m	metre
MAF	minor allele frequency
Met	methylation
mg	milligram
MgCl	magnesium chloride
min	minunte
Mio	million
miRNA	microRNA
ml	milliliter
mmol	millimole
MRI	magnetic resonance imaging
mRNA	messengerRNA
mU/l	microunits per milliliter
ng	nanogram
NGT	normal glucose tolerance
nl	nanoliter
N-TAD	N-terminal transactivation domain
O₂	oxygen
ODD	oxygen-dependent degradation domain
oGTT	oral glucose tolerance test
p300	E1A binding protein p300
p53	tumor protein p53
PAS domain	Per-Arndt-Sim domain
pg	picogram
pmol	picomole

pO₂	partial pressure of oxygen
PPAR	Peroxisome proliferator-activated receptor
Pro	proline
r	correlation coefficient
RNA	ribonucleic acid
RT-PCR	reverse transcription polymerase chain reaction
s	second
SAT	subcutaneous adipose tissue
sc	subcutaneous
SD	standard deviation
SEM	standard error of the mean
SFA	saturated fatty acids
SNP	single nucleotide polymorphism
SVF	stromal vascular fraction
T2D	type 2 diabetes
TG	triglycerides
TNFα	Tumor necrosis factor α
TSH	thyroid-stimulating hormone
VAT	visceral adipose tissue
VHL	Von-Hippel-Lindau tumor supressor protein
vis	visceral
vs.	versus
WHO	World Health Organisation
WHR	waist-to-hip ratio

1 Introduction

1.1 Obesity and its associated pathomechanisms

1.1.1 Obesity: a global health burden

Obesity constitutes an emerging global health problem. Worldwide 38% of adults aged 18 and over were overweight in 2014 (BMI 25-29,9kg/m²; 38% of men and 40% of women) and 13% were obese (BMI greater than or equal to 30kg/m²; 11% of men and 15% of women) (WHO). Obesity has more than doubled since 1980 (WHO). Figure 1 shows the world-wide prevalence of overweight by country in 2014.




Figure 1. Prevalence in % of overweight in adults 18 years and older by country in 2014
(http://gamapserver.who.int/gho/interactive_charts/ncd/risk_factors/overweight/flash/atlas.html).
Source: WHO.

Overweight and obesity used to be considered high-income country problems. However, they now depict a growing problem also in low- and middle-income countries, where the rate of increase of childhood overweight and obesity has recently been more than 30% higher than that of developed countries (WHO, 2014). Overweight and obesity are defined as abnormal or excessive fat accumulation that may impair health (WHO). The Body-Mass-Index (BMI) is a commonly used measure to quantitate obesity. It is defined as a person's weight in kilograms divided by the square of his height in meters (kg/m²). According to the WHO definition, a BMI greater than or equal 25 is overweight and a BMI greater than or equal 30 is obese. Factors determining BMI are mostly energy metabolism, nutrition and exercise, environmental factors and genetic factors (1).

Obesity significantly increases the risk of developing type 2 diabetes, hypertension, coronary heart disease, stroke, fatty liver disease, dementia, obstructive sleep apnea and some types of cancer (2–9). Globally, 44% of diabetes, 23% of ischemic heart disease and 7–41% of certain cancers are attributable to overweight and obesity (WHO, 2014). However, obesity itself is not alone relatable to these secondary diseases (10–13). Particular distinctions in body composition, fat distribution and adipose tissue function are considered to be more relevant for the development of obesity-associated disorders (14, 10, 11, 15), as individuals with low subcutaneous but increased visceral fat mass suffer an increased cardiometabolic risk (16, 17, 15, 13).

1.1.2 Adipose tissue

Adipose tissue is composed of different cells, connective tissue matrix, vascular and neural tissues (18). The cell population of adipose tissue consists of adipocytes and of a cell population called the stromal vascular fraction which includes monocytes, macrophages, pericytes, endothelial precursor cells, pre-adipocytes and adipocytes-derived mesenchymal stem cells (19). Adipocytes constitute the main cell population of adipose tissue. Free fatty acids and glycerol are stored in adipocytes as triglycerides (18). In periods of starvation these triglycerides can be mobilized for lipolysis and β -oxidation, thus serving as an energy storage system (18). Excess energy intake with limited energy expenditure conditions the accumulation of adipose tissue (18). In addition to the storage and release of lipids, adipose tissue secretes bioactive factors, adipokines, which have various functions concerning adipose tissue (e.g. regulating adipogenesis, immune cell migration in adipose tissue and adipocyte metabolisms) and the whole body (e.g. affection of the immune response, inflammation, glucose metabolism, insulin sensitivity and blood pressure) (14, 20, 21). Leptin, one of the most well studied adipokines, plays an important role in the regulation of satiety, appetite and energy expenditure (22). It lowers appetite in response to food intake by influencing signaling in the hypothalamus (22).

Adipose tissue develops from pluripotent mesenchymal stem cells which differentiate to preadipocytes and later to adipocytes (23). Key players involved in adipogenesis are the nuclear hormone receptor PPARG and CCAAT/enhancer binding protein α (C/EBP α). C/EBP α is a transcription factor that regulates PPARG (24). PPARG regulates transcriptional gene activation and is an essential regulator of adipocyte differentiation and lipid storage in AT (25).

About 80% of all body fat is located subcutaneously (SAT), mainly in the femorogluteal regions, back and anterior abdominal wall (18, 26). 10-20% of body fat in men and 5-8% in women are located in visceral depots (VAT) (18, 26). VAT is considered ectopic storage of adipose tissue and is mainly to be found in the central torso area surrounding organs. Ectopic fat storage further includes omental, pericardial, perirenal, and retroperitoneal fat deposition. Intraabdominal visceral accumulation of fat is associated with a much greater risk for the development of obesity associated comorbidities than subcutaneous fat accumulation (11, 15, 13). VAT and SAT differ biologically as VAT displays decreased insulin sensitivity, lower angiogenic potential, increased lipolytic activity, a different cellular composition and a different expression pattern of genes regulating adipocyte function (26). Whereas VAT is more vascular, innervated and contains a higher number of inflammatory and immune cells, SAT has a higher preadipocyte differentiating capacity and a lower percentage of large adipocytes (18). VAT drains into the portal vein, exposing the liver to FFA and adipokines released from adipocytes (14, 18). Adipokines activate hepatic immune mechanisms and stimulate the production of inflammatory mediators (27, 28). All of this could contribute to the deleterious consequences of visceral obesity, playing important roles in the development of obesity-associated metabolic and cardiovascular diseases.

The distribution of adipose tissue depends on various factors, like genetics, age, gender, total body fat content and energy balance (26). In young females, the subcutaneous abdominal fat area tends to be dominant over the abdominal visceral fat area. At the age of about 60, postmenopausal, this fat distribution shifts towards the predominantly visceral fat accumulation commonly found in men (29), due to the effect of sex hormones on fat distribution (30). The amount of visceral fat increases with age in both genders (29). A positive energy balance has only negligible impact on visceral fat accumulation. Hyper-energetic nutrition in monozygotic twins was associated with an increase in subcutaneous fat mass, whereas the variation in visceral fat did not exceed 10% (31). In contrast, body weight loss results in an over-proportional reduction of the visceral fat mass, probably due to the higher lipolytic activity of visceral compared with subcutaneous adipose tissue (32, 26).

1.1.3 Development of hypoxia in adipose tissue

Hypoxia is commonly found in obese adipose tissue. Several mechanisms are suspected to contribute to the development of hypoxia in obese adipose tissue (33). First, despite the

expansion of adipose tissue and total body fat mass in obesity, the proportion of the cardiac output and the dimension of blood flow to adipose tissue does not show an increase (34, 35). Second, whereas lean subjects show a postprandial increase in blood flow to adipose tissue, this rise cannot be observed in obese patients (36, 37). Third, the obesity-associated enlargement of adipocytes which can be up to 150-200 μ m in obese (38) can surmount the diffusion distance of oxygen of 100-200 μ m (39). Further, it has been shown that capillary density is lower in AT of obese individuals than in lean individuals (40, 41). Adipocyte hypertrophy results in adipose tissue expansion and, expanding the angiogenic potential, can be accompanied by insufficient vascularization (40). This results in increased distances between single adipocytes and capillaries, forming areas of adipose tissue with decreased blood perfusion and conditions the development of areas of adipose tissue hypoxia (40).

Hypoxia in adipose tissue could be verified through direct measurement of oxygen tension in adipose depots of obese mice with fiber-optic O₂ sensors (42, 43). A pO₂ of 48mmHg was measured in AT of lean mice, whereas for obese mice, the pO₂ was 15.2mmHg, providing direct evidence of hypoxia in obese adipose tissue in rodents (43). In humans, relative hypoxia in adipose tissue in obesity could be confirmed through a study showing that the pO₂ in subcutaneous AT of the upper arm is ~20mmHg lower in obese than in lean subjects (35).

1.1.4 Inflammation in adipose tissue

Adipose tissue expansion is closely related to chronic inflammation in adipose tissue, liver and skeletal muscle (44, 45, 33, 43). Chronic inflammation in adipose tissue gradually leads to the establishment of secondary pathological reactions such as insulin resistance, hyperinsulinemia and glucose intolerance (46). The relationship between obesity and inflammation was discovered with the observation that TNF α was elevated in obese adipose tissue and that inhibition of this cytokine improved glucose tolerance and insulin sensitivity (47). Further studies revealed that a key mechanism contributing to the establishment of obesity-induced inflammation is the accumulation of increased numbers of tissue macrophages (48, 49). These adipose tissue macrophages secrete different proinflammatory cytokines, like TNF α , IL1 β and IL6, which can directly inhibit insulin action in insulin target cells (i.e. hepatocytes, myocytes, adipocytes) (50, 51) or leak into the systemic circulation and cause insulin resistance through endocrine effects, further contributing to the establishment of an inflammatory state.

It is suggested that adipose tissue hypoxia triggers the inflammatory process (42, 43). However, the early events that initiate the cascade remain to be poorly understood. Part of this inflammation process is triggered by HIF1 α , a hypoxia inducible transcription factor that shows increased expression levels early in response to high fat diet fed mice (52). The state of relative adipocyte hypoxia provides the stimulus for HIF1 α induction (53, 54). HIF1 α can play a key initiating role in adipose tissue macrophage accumulation and inflammation by stimulating the production of adipocyte-derived chemokines and attracting macrophages (55, 52), thereby contributing to the establishment of an inflammatory state.

1.2 The transcription factor family of “hypoxia inducible factor”

1.2.1 The function of hypoxia inducible factors

Hypoxia-inducible factors are transcription factors that largely mediate hypoxia response (53). They are heterodimeric transcription factors consisting of an oxygen-labile α -subunit and a constitutively expressed β -subunit. Upon binding to specific sequences named hypoxia responsive elements (HREs) in their target genes, they mediate transcription of a large number of genes whose products are involved in erythropoiesis, angiogenesis, metabolic reprogramming, cell-cycle-regulation, and tumor progression (56, 57). Three different isoforms of the α -subunit, HIF1 α , HIF2 α and HIF3 α , exist and allow the formation of transcription factors with different functions upon dimerizing with HIF β . HIF1 α and HIF2 α are the most structurally similar, containing a basal Helix-loop-helix (bHLH) domain, two PAS domains, an oxygen-dependent-degradation domain (ODD), an N-terminal transactivation domain (N-TAD), and a C-TAD, of which the N-TAD partially overlaps with the ODD (58, 59). In normoxia, they are hydroxylated in at least one of the two critical prolines in their ODD, Pro402 or Pro564, allowing the E3 ubiquitin ligase von-Hippel-Lindau-Protein to ubiquitinate a leucine-residue, thereby marking them for proteasomal degradation (60, 61). In humans, this hydroxylation is catalyzed by three oxygen dependent HIF prolyl 4-hydroxylases (HIF-P4Hs 1-3) (62–65). Furthermore, the hydroxylase factor inhibiting HIF (FIH) hydroxylates an asparagine in the C-terminal transactivation domain (C-TAD) of HIF1 α and HIF2 α , preventing the binding of coactivator creb binding protein CBP/p300 and thereby inhibiting the full transcriptional activity of HIF1 α and HIF2 α in normoxia (66–68). In hypoxia, HIF-P4Hs and FIH are inhibited, resulting in reduced hydroxylation of the ODD and the CTAD, which prohibits degradation of HIF1 α and HIF2 α .

The stabilized HIF α -subunit is able to dimerize with the HIF β subunit, allowing the complex to bind to the hypoxia-responsive elements in the promoter region of its target genes (69, 70). These HIF1 α and HIF2 α target genes allow regulation of various biological processes such as erythropoiesis, angiogenesis, metabolic reprogramming, cell-cycle-regulation, and tumor progression, displaying master regulators of the transcriptional response to hypoxia (56, 57). A schematic presentation of HIF regulation under normoxic and hypoxic conditions is shown in Fig.2.




Figure 2. HIF regulation under normoxic and hypoxic conditions. Under normoxic conditions, HIF α is hydroxylated and ubiquitinated and thereby marked for proteasomal degradation. Under hypoxic conditions, HIF α is stabilized and forms a complex with HIF β and p300 that is capable of activating transcription of target genes. VHL = von-Hippel-Lindau-Protein, HRE = hypoxia responsive elements, Ub = ubiquitin, OH = hydroxyl group, Pro = proline, Asn = asparagin. (own graphic)

1.2.2 The role of hypoxia inducible factor 3 α

HIF3 α structurally differs from HIF1 α /2 α , possessing a leucine zipper domain (LZIP) as well as an LXXLL motif (71). HIF3 α lacks the CTAD, and thus cannot bind CBP/p300 (71, 72). The HIF3 α ODD differs from those of HIF1 α and HIF2 α as it contains only one hydroxylatable proline corresponding to Pro564 in HIF1 α , being oxygen-dependently regulated through hydroxylation by the HIF-P4Hs (64, 73). The function of HIF3 α differs from that of HIF1 α /2 α , displaying various roles in different biological processes. Multiple isoforms of HIF3 α exist, *HIF3A* being subject to extensive alternative splicing and different promoter- and transcription start site utilization (72, 74, 75). The existence of the variants HIF3 α -1, HIF3 α -2, HIF3 α -4, HIF3 α -7, HIF3 α -8, HIF3 α -9 and HIF3 α -10 could be

confirmed in various human tissues (76). A schematic representation of HIF1 α , HIF2 α and the HIF3 α isoforms is shown in figure 3.




Figure 3. Schematic presentation of HIF1 α , HIF2 α and the HIF3 α variant proteins. bHLH = basic helix-loop-helix, PAS = Per/Arndt/Sim, ODD = oxygen-dependent degradation domain, NTAD and CTAD = N- and C-terminal transactivation domain, LZIP = leucine zipper, P = prolines hydroxylated in the ODD, aa = amino acids (own graphic modified from (77)).

Some of these HIF3 α isoforms show weak or no transcriptional activity upon testing *in vitro* in human/mammalian cells and some of these isoforms even lack any TAD (71, 72, 76). The splice variant HIF3 α -4 was even able to suppress HIF1 α and HIF2 α -induced reporter construct expression in cultured cells in overexpression experiments (73). Overexpression of all the *HIF3A* variants in cell cultures was shown to inhibit the transcriptional activation of an HRE-reporter by HIF1 α and HIF2 α (77). These studies led to the conclusion that HIF3 α functions as a negative regulator of HIF1 α and HIF2 α . However, this view is derived from overexpression experiments performed in cell culture systems that may differ from the functioning of the endogenous protein *in vivo*. Recent studies reported that HIF3 α functions as a transcriptional activator in the hypoxia response in zebrafish embryos (78). According to this study, HIF3 α is capable of activating certain target genes independent or in collaboration with HIF1 α , suggesting a role of HIF3 α in glucose and amino acid metabolism, apoptosis, proteolysis, p53 signaling and PPAR signaling. HIF3 α has further been shown to be involved in adipocyte differentiation, as it was found to be induced during 3T3-L1 adipocyte differentiation and to be expressed in the differentiated adipocytes (79). Ectopic expression of *HIF3A* was shown to induce the expression of several genes related to adipocytes and to enhance adipogenic potential, suggesting HIF3 α to function as an

accelerator of adipogenesis (79). HIF3 α was further shown to be induced by insulin and 2-deoxy-glucose in rats, suggesting the transcriptional up-regulation of *HIF3A* to constitute a typical response to hypoglycaemia and glucoprivation *in vivo* (79, 80). Collectively, these studies suggest HIF3 α to be involved in metabolic derangements and in mechanisms resulting in the establishment of acquired obesity.

1.3 Genetics and epigenetics

1.3.1 Epigenetic mechanisms

Epigenetics is defined as an alteration in genomic signaling caused by modifications of chromosomes without affecting the genomic DNA sequence (81). This is achieved through chemical modification of the DNA region itself or through modifications of proteins that are closely associated with the locus (82). The specific targeting of genetic loci ensures a local effect of the alteration (82). Mechanisms allowing these alterations are histone modification (83, 84), DNA methylation (85), and long noncoding RNAs (86). Together, these mechanisms influence gene expression by adapting regions of the genome to achieve gene silencing or activation. Histone modification and DNA methylation interfere with the confirmation of chromatin, which influences the gene activity by altering the DNA accessibility to the transcription machinery (82). Figure 4 shows the most well investigated epigenetic mechanisms: DNA methylation and post-translational modifications of histone proteins such as acetylation, methylation, ubiquitination or phosphorylation.




Figure 4. Epigenetic mechanisms affecting the regulation of gene expression. Methylation of cytosine residues (C) within the DNA and modification of lysine (K) and arginine (R) residues of histone proteins. Me = methyl group, Ac = acetyl group, P = phosphor, Ub = ubiquitin. Source: <http://www.promega.de/resources/product-guides-and-selectors/protocols-and-applications-guide/epigenetics>

1.3.2 The methylation of DNA

DNA methylation is the predominant epigenetic modification in vertebrates (87). A methyl group from S-adenosine methionine is attached to the 5' position of a cytosine residue within a cytosine-phosphate-guanine (CpG) dinucleotide, creating 5-methylcytosine. The DNA methyltransferases DNMT3a, DNMT3b and DNMT3l catalyze *de novo* methylation, whereas DNMT1 is able to maintain methylation during DNA replication by recognizing hemimethylated CpGs and copying the DNA methylation pattern from the parental to the newly synthesized DNA strand (88, 89). This allows the patterns of DNA methylation to be relatively stable and to be maintained through cell generations. In addition to this, active and passive demethylation of DNA can allow dynamic regulation of gene expression (90). CpG dinucleotides are often located in short (approximately 1 kb) CpG-rich regions known as CpG-islands, which can be found in more than 50% of the genes in vertebrate genomes (91) and are often located within or close to the promoter region of genes. Methylation of CpG islands in proximity to the promoter results in repressed gene expression as the methylated DNA becomes inaccessible for the transcription machinery (92, 91). Furthermore, long-term silencing of genes such as imprinted genes, or tissue-specific genes in tissues where the gene is not expressed can be mediated via methylation of transcription start sites (91). Low methylation levels at the transcription start sites are, in contrast, associated with transcriptionally active genes. Contrary to this, DNA methylation in the gene body seems to be beneficial for transcription elongation (91). Thus, the effect of DNA methylation may not be causal, but have different functions depending on the context. DNA methylation is further involved in transcriptional regulation of genes and miRNA, control of alternative promoter usage, and alternative splicing (93–95). DNA transcription and phenotypic variation can be influenced by methylation. However, the determinants of DNA methylation itself remain poorly understood.

1.3.3 Epigenetics in association with obesity

Obesity is a multifactorial disease caused by an interaction of effects of genetics, epigenetics and the environment (96). Environmental factors depict dietary and physical activity patterns. More than 40 genetic variants are known to contribute to the etiology of obesity and more than 30 SNPs are associated with variations in BMI (97, 98). However, genetic variants identified so far are not sufficient in explaining the heritability of obesity. Therefore, epigenetic modifications, which affect long-term gene expression need to be acknowledged

as possible factors influencing the BMI (99). DNA methylation can be affected by both genetic and environmental factors (100, 101). It is plausible that an epigenetic change caused by integrating these factors results in an altered downstream functional outcome, influencing the phenotype (102). This might possibly play a role in the pathophysiology of obesity and adverse fat distribution. Previous studies confirmed functionally relevant methylation differences between different adipose depots in pigs, supporting the role of epigenetics in the regulation of fat distribution (103). Genome-wide analyses of methylation at CpG sites in relation to BMI revealed different sites in *HIF3A* to show an association between methylation and BMI in whole blood and in adipose tissue (104–106).

1.3.4 Single nucleotide polymorphisms in obesity

Single nucleotide polymorphisms (SNPs) are variations at specific locations in the genome. These variations consist of sequence differences in single nucleotide positions (107). In every 1000 bases along the human chromosomes, approximately one nucleotide position differs between any two copies of that chromosome (108, 109). The locus is regarded as polymorphic when it exists in two or more variants and the allele frequency of the most common variant is <90% (110). Some of these SNPs directly affect protein structure or expression levels of genes. Therefore, they directly influence the transcriptome, and can, if they are located in relevant sites, represent genetic mechanisms in the etiology of various diseases. To date, more than 30 SNPs are known to be associated with BMI, which together explain about 1.5% of interindividual variation in BMI (98). DNA sequence variation can further influence DNA methylation levels (111, 112). Two SNPs have been shown to be associated with methylation at the sites mentioned above. However, they did not show an association with BMI (105).

1.4 Main hypothesis

Here we tested the hypothesis that *HIF3A* mRNA expression and CpG-sites methylation in AT is related to parameters of adipose tissue distribution and function. In paired samples of subcutaneous AT (SAT) and visceral AT (VAT) from 603 individuals with a wide range of age, BMI and obesity-related parameters, we investigated whether *HIF3A* mRNA expression is fat depot-specific, altered in obesity and related to measures of AT function and insulin sensitivity. In a subgroup of 548 individuals, we investigated the effects on *HIF3A* genetic variants on *HIF3A* AT expression and *HIF3A* methylation of CpG-sites.

1.5 Aim of this study

The aim of our study is to further investigate the relation between hypoxia inducible factors and obesity. We thereby aim to contribute to a better understanding of the pathophysiological mechanisms involved in obesity. Genome-wide analysis of methylation at CpG sites in relation to BMI revealed several sites of *HIF3A* to show an association between methylation and BMI in whole blood and in adipose tissue (104–106). Two SNPs have been shown to be associated with methylation at these sites, but to be independent of BMI (105). Downstream effects of obesity-induced epigenetic changes resulting in modified signaling of the HIF-system could mediate important mechanisms in answer to energetic dysbalance. It is therefore of great importance to further investigate the coherence between hypoxia inducible factors and development of obesity-associated comorbidities, possibly revealing important insights into pathophysiological processes concerning AT-inflammation, insulin resistance or the etiology of obesity. We therefore tested the hypothesis that expression of *HIF3A* in human subcutaneous and visceral AT is related to BMI, parameters of fat distribution and AT function, metabolic traits, genetic variation and methylation of CpG-sites in *HIF3A*.

2 Manuscript of the publication

SCIENTIFIC REPORTS

OPEN

Hypoxia-inducible factor 3A gene expression and methylation in adipose tissue is related to adipose tissue dysfunction

Received: 20 November 2015

Accepted: 26 May 2016

Published: 27 June 2016

Susanne Pfeiffer^{1,*}, Jacqueline Krüger^{2,*}, Anna Maierhofer³, Yvonne Böttcher², Nora Klötting^{1,2}, Nady El Hajj³, Dorit Schleinitz², Michael R. Schön⁴, Arne Dietrich^{2,5}, Mathias Fasshauer^{1,2}, Tobias Lohmann⁶, Miriam Dreßler⁶, Michael Stumvoll¹, Thomas Haaf³, Matthias Blüher¹ & Peter Kovacs²

Recently, a genome-wide analysis identified DNA methylation of the *HIF3A* (*hypoxia-inducible factor 3A*) as strongest correlate of BMI. Here we tested the hypothesis that *HIF3A* mRNA expression and CpG-sites methylation in adipose tissue (AT) and genetic variants in *HIF3A* are related to parameters of AT distribution and function. In paired samples of subcutaneous AT (SAT) and visceral AT (VAT) from 603 individuals, we measured *HIF3A* mRNA expression and analyzed its correlation with obesity and related traits. In subgroups of individuals, we investigated the effects on *HIF3A* genetic variants on its AT expression (N = 603) and methylation of CpG-sites (N = 87). *HIF3A* expression was significantly higher in SAT compared to VAT and correlated with obesity and parameters of AT dysfunction (including CRP and leucocytes count). *HIF3A* methylation at cg22891070 was significantly higher in VAT compared to SAT and correlated with BMI, abdominal SAT and VAT area. Rs8102595 showed a nominal significant association with AT *HIF3A* methylation levels as well as with obesity and fat distribution. *HIF3A* expression and methylation in AT are fat depot specific, related to obesity and AT dysfunction. Our data support the hypothesis that HIF pathways may play an important role in the development of AT dysfunction in obesity.

Obesity and its associated comorbidities constitute an evolving health burden worldwide¹. Obesity is closely related to chronic inflammation in adipose tissue, liver and skeletal muscle², which may contribute to chronic systemic inflammation, insulin resistance, and deterioration in glucose and lipid metabolism³. Upon weight gain, adipocyte hypertrophy may lead to hypoxia in adipose tissue which is considered as a causative factor in adipose tissue dysfunction^{4–7}. It has been recently shown that adipose tissue expression of *hypoxia inducible factor (HIF) 1A* (*HIF1A*) increases in mice exposed to high fat diet⁴. In states of relative adipose tissue hypoxia, induction of HIF1 α ^{5,6} stimulates accumulation of macrophages in adipose tissue^{4,7} and the production of adipocyte-derived pro-inflammatory cytokines. HIFs are heterodimeric transcription factors that mediate hypoxia response in various tissues⁶. They consist of an oxygen-labile α -subunit and a constitutively expressed β -subunit. Three existing isoforms of the α -subunit, HIF1 α , HIF2 α and HIF3 α , allow the formation of transcription factors with different functions upon dimerizing with HIF β . Multiple isoforms of HIF3 α exist⁸. HIF3 α is capable of activating certain target genes independent or in collaboration with HIF1 α , suggesting a role of HIF3 α in glucose and amino acid metabolism, apoptosis, proteolysis, p53 signaling and PPAR signaling. In addition, HIF3 α has been shown to play a role in adipocyte differentiation^{9,10}.

¹Department of Medicine, Dermatology und Neurology, Department of Endocrinology und Nephrology, University of Leipzig, Leipzig, Germany. ²Leipzig University Medical Center, IFB AdiposityDiseases, University of Leipzig, Leipzig, Germany. ³Institute of Human Genetics, University of Würzburg, Würzburg, Germany. ⁴Clinic of Visceral Surgery, Städtisches Klinikum Karlsruhe, Karlsruhe, Germany. ⁵Department of Surgery, University of Leipzig, Leipzig, Germany. ⁶Municipal Clinic Dresden-Neustadt, Dresden, Germany. *These authors contributed equally to this work. Correspondence and requests for materials should be addressed to P.K. (email: peter.kovacs@medizin.uni-leipzig.de)

Recent genome-wide analysis of DNA methylation in whole blood and human adipose tissue revealed an association of methylation at three CpG sites in intron 1 of *HIF3A* with BMI^{11–13}. In addition, two single nucleotide polymorphisms (SNPs) rs8102595 and rs3826795, have been shown to be associated with methylation at these sites, yet to be independent of BMI¹¹. The strong relationship of *HIF3A* methylation and obesity was also shown in neonates¹⁴. Furthermore, gene-diet interactions between the methylation-associated SNP rs3826795 and vitamin B intake were recently reported, providing a potential causal link between the epigenetic status and obesity¹⁵.

Further investigation of the relationship between HIFs and development of obesity-associated comorbidities might reveal important insights in pathophysiological processes concerning AT inflammation and/or insulin resistance in the etiology of obesity related metabolic diseases. We therefore tested the hypothesis that expression of *HIF3A* in human subcutaneous and visceral adipose tissue is related to obesity, parameters of fat distribution and adipose tissue function. We further assessed the relationship between the AT expression, genetic variation (rs8102595 and rs3826795) and methylation of CpG-sites in *HIF3A*.

Material and Methods

Study participants. A total of 288 Caucasian men and 577 women were included in the study (Table 1). According to the ADA criteria, 343 subject were diagnosed with type 2 diabetes (T2D) and 484 had normal glucose tolerance (NGT)¹⁶. Paired samples of visceral adipose tissue and subcutaneous adipose tissue were obtained from 603 individuals following open abdominal surgery for gastric banding, cholecystectomy, weight reduction surgery, abdominal injuries or explorative laparotomy. Patients with end-stage malignant diseases were excluded from the study. All adipose tissue samples were frozen immediately in liquid nitrogen after explantation and stored at -80°C . Six-hundred and three subjects (mean age 50 ± 14 years, mean BMI $43.6 \pm 13.0 \text{ kg/m}^2$) were included into adipose tissue *HIF3A* mRNA expression analysis. DNA methylation analysis was performed in a subgroup of 87 subjects (mean age 58 ± 15 years, mean BMI $32.9 \pm 12.7 \text{ kg/m}^2$). Genotyping was done in 548 individuals overlapping with adipose tissue biopsy donors (mean age 50 ± 14 years, mean BMI $34.6 \pm 13.6 \text{ kg/m}^2$).

Phenotypic characterization including anthropometric measurements, body fat analysis (bioimpedance analyses or dual-energy X-ray absorptiometry) and metabolic parameters such as fasting plasma glucose and insulin, a 75-g oral glucose tolerance test (oGTT), HbA1c, lipoprotein-, triglyceride-, free fatty acid- and adipokine serum concentrations was performed as previously described^{17,18}. Measurement of abdominal visceral and subcutaneous fat areas ($N = 245$) was performed using computed tomography (CT) or MRI scans. All subjects had a stable weight, defined as the absence of fluctuations of $>2\%$ of body weight for at least 3 months before surgery. In addition, adipocytes and cells of the stromal vascular fraction were isolated from adipose tissue samples of 35 subjects (18 men, 17 women). Adipocytes were isolated by collagenase (1 mg/ml) digestion. To determine cell size distribution and adipocyte number, aliquots of adipocytes were fixed with osmic acid and counted in a Coulter counter as previously described¹⁹. The study was approved by the ethics committee of the University of Leipzig (approval number: 159-12-21052012) and all subjects gave written informed consent. All methods were carried out in accordance with the approved guidelines.

Analysis of human *HIF3A* mRNA expression. Briefly, human *HIF3A* mRNA expression was measured by qRT-PCR using TaqMan Gene Expression Assay (Applied Biosystems, Darmstadt, Germany). Total RNA was isolated from adipose tissue samples using the Qiacube System (Qiagen, Hilden, Germany), and $2 \mu\text{g}$ RNA were reverse transcribed with standard reagents (Life Technologies). Further details including PCR conditions are provided in the Supplemental material. The following Gene Expression Assay was used: Hs00541709_M1 (tagging the transcripts NM_022462.4, NM_152794.3, NM_152795.3, NM_152796.4). *HIF3A* mRNA expression was calculated relative to the mRNA expression of *hypoxanthine guanine phosphoribosyltransferase 1* (*HPRT1*), determined by the assay Hs01003267_M1 (Applied Biosystems, Darmstadt, Germany). Expression of *HIF3A* and *HPRT1* mRNA were quantified by using the second derivative maximum method of the TaqMan Software (Applied Biosystems).

For expression analysis of *HIF3A* in adipocytes and stromal vascular fraction, total RNA was isolated from adipocytes and stromal vascular fraction extracted from 35 paired samples of subcutaneous and visceral adipose tissue. 300 ng RNA were reverse transcribed with standard reagents and from each RT-PCR, $23.5 \mu\text{l}$ was amplified in a $40 \mu\text{l}$ PCR using the Taqman Gene Expression Assay and the TaqMan Fast Advanced Mastermix according to the manufacturer's instruction. *HIF3A* mRNA expression was calculated relative to the mRNA expression of *HPRT1* mRNA or *18S rRNA*, determined by the assay Hs01003267_m1 (Applied Biosystems, Darmstadt, Germany).

DNA extraction and bisulfite conversion. Briefly, genomic DNA was extracted using the DNeasy Blood and Tissue Kit (Qiagen, Hilden, Germany) and bisulfite conversion was performed using the Epitect Bisulfite Kit (Qiagen, Hilden, Germany) according to the manufacturer's protocol.

Determining CpG methylation levels. PCR and sequencing primers were designed using the PyroMark Assay Design 2.0 software (Qiagen, Hilden, Germany). DNA fragments were amplified from bisulfite-converted DNA using forward primer 5'-TGGTTGAAGGGTTATTAGGG-3' and biotinylated reverse primer 5'-ACTCTATCCCACCCCTTTT-3'. The PCR reaction mixture and cyclor program are provided in the Supplementary material. Bisulfite pyrosequencing was performed on a PyroMark Q96MD pyrosequencing system (Qiagen) using the PyroMark Gold Q96 CDT reagent kit (Qiagen) and the Pyro Q-CpG software (Qiagen). Percentage of methylation at eleven individual CpG sites within intron 1 of *HIF3A* were determined using three different sequencing primers (Assay 1: 5'-TTTAGGGGTGTAGG-3'; Assay 2: 5'-GGTGAGATGATTTTATAGGAA-3'; Assay 3: 5'-GTTAAGAGGGGTTTTTATT-3'). Assay 1 included seven CpGs, Assay 2 only one CpG and Assay 3 three CpGs. The sixth CpG site in Assay 1, the CpG site in Assay 2 and

	Total	Lean	Overweight	Obese	NGT	T2D	CRP < 5
N	865	88	73	704	484	343	347
Men/Women	288/577	40/48	36/37	212/492	155/329	124/219	129/218
Age (years)	49 ± 13	62 ± 15 ^{aaa}	62 ± 14	46 ± 11 ^{ccc}	47 ± 15 ^{ddd}	51 ± 10	49 ± 13
BMI (kg/m ²)	44.3 ± 12.6	22.1 ± 2.3 ^{aaa}	27.2 ± 1.3 ^{bbb}	48.9 ± 9.0 ^{ccc}	40.9 ± 12.9 ^{ddd}	49.4 ± 10.8	41.8 ± 11.7 ^{eee}
Body weight (kg)	129 ± 39	63 ± 9 ^{aaa}	78 ± 9 ^{bbb}	142 ± 30 ^{ccc}	119 ± 41 ^{ddd}	143 ± 33	122 ± 38 ^{eee}
Height (m)	1.6 ± 0.1	1.68 ± 0.09	1.69 ± 0.09	1.69 ± 0.09	1.69 ± 0.1	1.69 ± 0.1	1.70 ± 0.09
Waist (cm)	126.8 ± 26.2	77.0 ± 13.7 ^{aaa}	96.8 ± 13.8 ^{bbb}	137.1 ± 19.7 ^{ccc}	116.2 ± 28.2 ^{ddd}	142.6 ± 22.0	124.0 ± 27.9
Hip (cm)	129.1 ± 28.7	86.3 ± 9.7 ^{aaa}	102.4 ± 11.4 ^{bbb}	141.5 ± 21.0 ^{ccc}	126.0 ± 30.3 ^{ddd}	135.7 ± 24.7	127.0 ± 28.5
WHR	0.95 ± 0.13	0.90 ± 0.11 ^{aa}	0.94 ± 0.08	0.97 ± 0.14	0.91 ± 0.12 ^{ddd}	1.05 ± 0.08	0.95 ± 0.13
Visceral Fat area (cm ²)	240 ± 172	45 ± 29 ^{aaa}	119 ± 60 ^{bbb}	313 ± 154 ^{ccc}	177 ± 142 ^{ddd}	392 ± 150	209 ± 175
SC fat area (cm ²)	1093 ± 789	52 ± 26 ^{aaa}	273 ± 171 ^{bbb}	1509 ± 559 ^{ccc}	992 ± 817 ^{ddd}	1386 ± 671	920 ± 814 ^e
CT ratio (vis/sc)	0.4 ± 0.5	1.9 ± 0.9 ^{aaa}	0.5 ± 0.3 ^{bb}	0.2 ± 0.1 ^{ccc}	0.3 ± 0.4	0.5 ± 0.7	0.4 ± 0.5
Body Fat (%)	41.4 ± 11.5	19.0 ± 3.5	24.3 ± 3.9	45.1 ± 8.0	38.5 ± 13.0	44.3 ± 9.4	39.1 ± 11.0
CRP (mg/dl)	11.4 ± 14.4	14.9 ± 22.9	9.1 ± 13.1	11.2 ± 13.1	11.2 ± 15.6	12.1 ± 13.2	2.4 ± 1.5
IL-6 (pg/ml)	6.0 ± 5.2	2.2 ± 3.3	2.8 ± 2.4	7.0 ± 5.3	5.2 ± 4.6	7.5 ± 6.0	4.1 ± 1.3
HbA1c (%)	6.1 ± 1.2	5.3 ± 0.4	5.7 ± 0.6	6.1 ± 1.2	5.5 ± 0.5	6.9 ± 1.4	6.0 ± 1.0
oGTT2h (mmol/l)	7.0 ± 2.6	6.0 ± 1.0	6.1 ± 0.9	7.4 ± 2.9	6.3 ± 1.0	14.8 ± 5.9	6.6 ± 1.7
FPG (mmol/l)	6.5 ± 2.5	5.5 ± 1.0	5.9 ± 1.5	6.7 ± 2.7	5.4 ± 1.0	8.1 ± 3.2	6.1 ± 2.1
FPI (pmol/l)	123.1 ± 133.8	10.8 ± 20.6	68.0 ± 92.5	146.4 ± 137.2	62.4 ± 70.8	206.2 ± 156.4	109.7 ± 121.7
GIR (μmol/kg/min)	75.1 ± 33.4	102.5 ± 18.5	77.7 ± 25.8	56.9 ± 31.2	90.6 ± 21.4	30.5 ± 23.5	85.3 ± 28.3
Total cholesterol (mmol/l)	4.9 ± 1.0	5.1 ± 0.8	5.0 ± 1.1	4.9 ± 1.0	4.9 ± 1.0	4.9 ± 1.0	4.9 ± 1.0
HDL-C (mmol/l)	1.2 ± 0.3	1.7 ± 0.5	1.4 ± 0.3	1.1 ± 0.3	1.3 ± 0.4	1.1 ± 0.3	1.2 ± 0.4
LDL-C (mmol/l)	3.1 ± 0.9	2.8 ± 1.0	3.2 ± 0.8	3.1 ± 0.9	3.1 ± 0.9	3.0 ± 0.8	3.1 ± 0.9
FFA (mmol/l)	0.5 ± 0.4	0.2 ± 0.2	0.3 ± 0.3	0.6 ± 0.3	0.3 ± 0.3	0.8 ± 0.3	0.5 ± 0.3
TG (mmol/l)	1.8 ± 1.1	1.1 ± 0.4 ^{aaa}	1.2 ± 0.5	1.9 ± 1.1 ^{ccc}	1.4 ± 0.9 ^{ddd}	2.1 ± 1.1	1.8 ± 1.1
Leptin (ng/ml)	39.3 ± 24.2	4.8 ± 3.7 ^{aaa}	12.4 ± 7.0 ^{bbb}	45.4 ± 21.8 ^{ccc}	37.1 ± 23.9	41.2 ± 25.0	35.6 ± 22.7 ^e
Adiponectin (μg/ml)	6.9 ± 4.4	14.3 ± 6.2 ^{aaa}	8.8 ± 3.5 ^{bbb}	6.0 ± 3.3 ^{ccc}	8.5 ± 4.7 ^{ddd}	4.9 ± 3.1	7.1 ± 4.2
Albumin (g/L)	28.1 ± 18.9	32.9 ± 7.7	34.4 ± 13.2	26.9 ± 20.2	27.1 ± 18.8	28.6 ± 19.6	29.5 ± 20.0
ALAT (μkat/l)	0.6 ± 0.5	0.4 ± 0.3 ^{aaa}	0.5 ± 0.3	0.7 ± 0.5 ^{ccc}	0.6 ± 0.4 ^d	0.7 ± 0.5	0.7 ± 0.4
ASAT (μkat/l)	0.6 ± 2.3	0.4 ± 0.3	0.4 ± 0.2	0.6 ± 2.5	0.6 ± 3.0	0.6 ± 0.5	0.5 ± 0.3
gGT (μkat/l)	0.9 ± 1.3	1.0 ± 1.4	1.0 ± 1.5	0.8 ± 1.3	0.8 ± 1.0 ^d	1.0 ± 1.7	0.7 ± 0.9 ^e
TSH (mU/l)	1.9 ± 7.9	1.5 ± 2.1	1.5 ± 1.9	2.1 ± 8.8	1.7 ± 1.8	2.3 ± 12.4	1.4 ± 1.0
ft3 (pg/ml)	4.6 ± 0.9	4.5 ± 1.0	4.4 ± 0.7	4.7 ± 0.9 ^e	4.5 ± 0.9	4.7 ± 0.9	4.6 ± 0.9
ft4 (pmol/l)	17.1 ± 1.4	17.2 ± 3.3	17.5 ± 3.2	16.9 ± 3.5	16.9 ± 3.4 ^d	17.4 ± 3.4	17.6 ± 3.3 ^e
Leucocytes/nl	8.1 ± 2.7	7.5 ± 3.2	7.4 ± 3.1	8.2 ± 2.6 ^e	2.1 ± 0.5	2.1 ± 0.5	7.4 ± 2.2 ^e
Erythrocytes (Mio/μl)	4.7 ± 0.8	4.6 ± 2.7	4.3 ± 0.9	4.7 ± 0.4	4.7 ± 1.0	4.7 ± 0.4	4.7 ± 0.4
Thrombocytes (10 ⁹ /l)	260 ± 81	252 ± 108	241 ± 71	261 ± 79	261 ± 73	259 ± 88	232 ± 66 ^e

Table 1. Anthropometric and metabolic characteristics of study participants. Data are means ± SD; ^{a,b,c,d,e}_p < 0.05, ^{aa,bb,cc,dd,ee}_p < 0.01, ^{aaa,bbb,ccc,ddd,eee}_p < 0.001 for comparison between (a) lean and obese, (b) lean and overweight, (c) overweight and obese, (d) type 2 diabetes subjects (T2D) and subjects with normal glucose tolerance (NGT) and (e) CRP < 5 and the entire cohort. 51 subjects with type 1 diabetes or impaired glucose tolerance were not considered for group comparison. BMI – Body Mass Index, WHR – waist-to-hip ratio, sc – subcutaneous, TG – Triglycerides, ALAT – alanine aminotransferase, ASAT – aspartate aminotransferase, gGT – Gamma-glutamyl transferase, TSH – thyroid-stimulating hormone, ft3 – free triiodothyronine, ft4 – free tetraiodothyronine, CRP – C-reactive protein, IL-6 – Interleukin 6, HbA1c – Glycohemoglobin, oGTT – oral Glucose Tolerance Test, FPG – Fasting plasma glucose, FPI – Fasting plasma insulin, GIR – Glucose infusion rate during the steady state of an euglycemic hyperinsulinemic clamp, HDL-C – high Density Lipoprotein Cholesterol, LDL-C – Low Density Lipoprotein Cholesterol, FFA – Free Fatty Acids.

the third CpG site in Assay 3 correspond to the CpG sites on the 450 K array reported elsewhere¹¹. In our experience, the average methylation difference between technical replicates is approximately one percentage point.

Genotyping of *HIF3A* SNPs. Genomic DNA was extracted from blood using the Quick Gene DNA whole blood Kit (Kurabo, Japan). Genotyping of the two previously reported SNPs rs8102595 (A/G) and rs3826795 (G/A)¹¹ was performed using the TaqMan SNP Genotyping assay (Applied Biosystems; C_29247492_10; C_31640839_10). To assess genotyping reproducibility, a random ~5% selection of the sample were re-genotyped for all SNPs; all genotypes matched initial designated genotypes. Potential functional significance of the studied genetic variants was checked using the Regulome Database, which includes public datasets from GEO, the ENCODE project, and published literature²⁰.




Figure 1. *HIF3A* mRNA expression in human subcutaneous (SAT, n = 584) and visceral (VAT, n = 588) adipose tissue. In the entire study cohort (A), but also in subgroups of men (SAT, n = 108; VAT, n = 110) and women (SAT, n = 231; VAT, n = 230) (Subjects with T2D were excluded from analysis) (B). Expression of *HIF3A* is significantly higher in subcutaneous (SAT) compared to visceral (VAT) adipose tissue. (C) *HIF3A* mRNA expression in adipocytes (n = 35) and cells of the stromal vascular fraction (SVF). *HIF3A* is significantly higher expressed in adipocytes compared to cells of the SVF in both compartments (D) *HIF3A* mRNA expression in relation to adipocyte cell size in subcutaneous (SAT) and visceral (VAT) adipose tissue. Individuals were categorized by mean SAT and VAT adipocyte size into deciles. Comparison of *HIF3A* mRNA expression between individuals with the lowest versus highest mean adipocyte size decile reveals that *HIF3A* is more highly expressed in subjects with higher mean adipocyte volume. Data are presented as means ± SEM. **p < 0.01, ***p < 0.001, AU-arbitrary units.

Statistical Analyses. All non-normally distributed parameters were logarithmically transformed to approximate a normal distribution. To analyze differences in *HIF3A* methylation/expression levels between visceral and subcutaneous adipose tissue, paired two-tailed t-tests were applied. To test for group differences (e.g. lean vs. obese, NGT vs. T2D) two tailed t-tests were used. Pearson's correlation coefficients were used to assess bivariate correlation with phenotypes related to obesity, fat distribution and glucose and insulin homeostasis. Linear regression models were used to control for confounders such as age, gender and BMI. To test SNPs for genetic associations with mRNA expression, DNA methylation and metabolic traits, linear regression analysis adjusted for respective covariates was applied. Association studies on type 2 diabetes (T2D) and obesity (lean with BMI < 25 kg/m² vs. obese with BMI ≥ 30 kg/m²) were done using logistic regression analyses. P-values ≤ 0.05 were considered to provide nominal evidence for association. Two-sided p-values are reported without adjustments for multiple testing. The analysis of associations with quantitative traits was restricted to nondiabetic subjects to avoid diabetes status or treatment masking potential effects of the variants on these parameters. Statistical analyses were performed using SPSS statistics version 20.0.1 (SPSS, Inc., Chicago, IL, USA).

Results

***HIF3A* mRNA expression is fat depot related.** Analysis of paired subcutaneous and visceral adipose tissue samples revealed significantly higher *HIF3A* mRNA expression in subcutaneous compared to visceral adipose tissue (Fig. 1A). The fat depot differences in *HIF3A* expression could be confirmed in both genders (Fig. 1B). There was no significant difference in both subcutaneous and visceral adipose tissue *HIF3A* mRNA expression between individuals with normal glucose tolerance (NGT) and with type 2 diabetes (Supplementary Figure).

We further analyzed the contribution of adipocytes and stromal vascular fraction cells on whole adipose tissue *HIF3A* mRNA expression. Analysis of visceral and subcutaneous stromal vascular fraction showed significantly higher *HIF3A* mRNA levels in subcutaneous compared to visceral stromal vascular fraction (p < 0.05) (subcutaneous 0.56 ± 0.84 and visceral 0.37 ± 0.57). In paired samples of adipocytes and stromal vascular fraction cells we found significantly higher *HIF3A* expression in adipocytes compared to stromal vascular fraction cells both in subcutaneous and visceral fat compartments (Fig. 1C). There was no significant fat depot-related difference in *HIF3A* mRNA expression of isolated adipocytes. We further sought to determine *HIF3A* mRNA expression

	HIF3A mRNA Expression in subcutaneous adipose tissue			HIF3A mRNA Expression in visceral adipose tissue		
	r	p-value	adj. p-value	r	p-value	adj. p-value
Age (years)	−0.23	4.61×10^{-5}	0.032	−0.237	3.08×10^{-5}	0.076
BMI (kg/m ²)	0.239	2.86×10^{-5}	0.017^a	0.283	5.46×10^{-7}	8.84×10^{-4a}
Body weight (kg)	0.235	5.56×10^{-5}	0.467 ^a	0.263	5.45×10^{-6}	0.280 ^a
Height (m)	0.044	0.458	0.467	0.001	0.983	0.538
Waist (cm)	0.472	8.41×10^{-9}	0.010	0.515	1.89×10^{-10}	0.048
Hip (cm)	0.387	2.13×10^{-5}	0.425	0.442	6.73×10^{-7}	0.628
WHR	0.172	0.067	0.018	0.139	0.135	0.033
Visceral fat area (cm ²)	0.391	3.19×10^{-5}	0.636	0.442	1.71×10^{-6}	0.479
SC fat area (cm ²)	0.392	2.99×10^{-5}	0.240	0.465	4.06×10^{-7}	0.604
CT ratio (sc/vis)	−0.259	7.04×10^{-3}	0.165	−0.319	7.80×10^{-4}	0.325
Body fat (%)	0.324	0.017	0.055 ^a	0.442	8.23×10^{-4}	0.013^a
CRP (mg/dl)	−0.138	0.021	1.8×10^{-3}	−0.153	0.010	3.19×10^{-4}
Leucocytes/nl	−0.127	0.032	3.05×10^{-3}	−0.133	0.024	1.13×10^{-3}
Met Blood (%)	0.054	0.720	0.618	0.023	0.876	0.772
Met SAT (%)	−0.054	0.687	0.482	−0.088	0.498	0.345
Met VAT (%)	0.060	0.648	0.667	−0.045	0.729	0.757
Leptin mRNA sc	0.227	2.82×10^{-4}	2.37×10^{-4}	0.216	5.28×10^{-4}	1.41×10^{-3}
Leptin mRNA vis	0.117	0.063	0.043	0.195	2.0×10^{-3}	2.60×10^{-3}
PPARG mRNA sc	0.001	0.977	0.939	0.002	0.967	0.878
PPARG mRNA vis	0.050	0.253	0.439	0.111	0.010	0.023

Table 2. Correlation analyses of subcutaneous and visceral adipose tissue *HIF3A* mRNA expression with metabolic parameters, methylation levels and mRNA expression of *leptin* and *PPARG*. r - correlation coefficient (Pearson adj. - p-value adjusted to age, sex and BMI), ^aadjusted for sex and age; BMI - Body Mass Index, WHR - waist-to-hip ratio, sc - subcutaneous, CRP - C-reactive protein, Met Blood (%) / Met SAT (%) / Met VAT (%) - Methylation of cg22891070 in *HIF3A* in blood/SAT/VAT.




Figure 2. *HIF3A* mRNA expression in subcutaneous and visceral adipose tissue in relation to CRP serum concentration categories (n = 318) and leucocyte counts (n = 326). A significant inverse relationship between both CRP level (A) and leucocyte count (B) and expression of *HIF3A* in both compartments can be observed. Data are presented as means \pm SEM. ***p < 0.001, AU-arbitrary units.

in relation to adipocyte cell size. Comparison of *HIF3A* mRNA expression between individuals with the lowest versus highest mean adipocyte size decile reveals that *HIF3A* is more highly expressed in subjects with higher mean adipocyte volume (Fig. 1D).

***HIF3A* mRNA expression in adipose tissue correlates with parameters of obesity, systemic inflammation, glucose metabolism and mRNA expression of genes regulating adipogenesis (*leptin*, *PPARG*).** *HIF3A* mRNA expression in visceral and subcutaneous adipose tissue correlated significantly with BMI, body weight, waist and hip circumferences, abdominal visceral and subcutaneous fat area, %body fat, free fatty acid, triglyceride, alanine aminotransferase (ALAT), leptin serum concentrations and with the mRNA expression of *Leptin* (Table 2 and Supplementary Table 1). Furthermore, there were significant inverse correlations between subcutaneous and visceral adipose tissue *HIF3A* expression and age, CT ratio, adiponectin and C-reactive protein (CRP) serum concentrations (Fig. 2A) and leucocyte count (Fig. 2B). Only in visceral adipose tissue, *HIF3A* expression correlated with fasting plasma insulin, thyroid-stimulating hormone (TSH),

	rs8102595			rs3826795		
	A/A	A/G + G/G	p-value	A/A+ A/G	G/G	p-value
N	446	95		208	336	
Men/Women	151/295	32/63		73/135	110/226	
Age	52.83 ± 15.79	55.48 ± 15.44	0.482	49.56 ± 15.31	50.72 ± 14.69	0.278
BMI (kg/m ²)	43.48 ± 13.74	42.51 ± 13.50	0.239	43.64 ± 14.04	42.93 ± 13.32	0.908
Body weight (kg)	126.86 ± 42.81	124.57 ± 40.14	0.680	128.42 ± 45.54	124.60 ± 41.15	0.769
Height (m)	1.69 ± 0.09	1.69 ± 0.9	0.628	1.69 ± 0.09	1.69 ± 0.09	0.763
Waist (cm)	124.26 ± 29.98	121.84 ± 30.09	0.798	124.46 ± 30.43	122.85 ± 29.87	0.935
Hip (cm)	130.53 ± 28.99	128.59 ± 28.38	0.851	129.54 ± 28.38	130.08 ± 29.56	0.676
WHR	0.95 ± 0.13	0.96 ± 0.16	0.316	0.96 ± 0.16	0.94 ± 0.12	0.921
VAT area (cm ²)	242.93 ± 173.84	237.02 ± 159.92	0.575	256.05 ± 183.40	228.98 ± 159.97	0.674
SAT area (cm ²)	1095.74 ± 795.48	1129.73 ± 819.78	0.536	1122.85 ± 774.80	1094.46 ± 817.64	0.902
VAT mean	123.00 ± 20.82	122.08 ± 20.60	0.999	119.69 ± 25.71	124.66 ± 17.25	0.014
SAT mean	127.37 ± 19.89	127.51 ± 17.42	0.486	126.50 ± 19.04	127.99 ± 19.84	0.334
VAT max	209.23 ± 58.51	230.21 ± 96.06	0.060	210.73 ± 74.84	213.66 ± 63.47	0.109
SAT max	214.28 ± 70.88	249.22 ± 110.69	1.23 × 10⁻³	224.71 ± 80.22	217.94 ± 79.94	0.987
CT ratio (vis/sc)	0.47 ± 0.63	0.38 ± 0.30	0.922	0.40 ± 0.42	0.48 ± 0.66	0.826
Body fat (%)	41.95 ± 11.35	42.26 ± 11.72	0.496	41.15 ± 11.88	42.57 ± 11.11	0.607
CRP (mg/dl)	12.04 ± 15.09	11.20 ± 16.05	0.935	13.09 ± 15.67	11.34 ± 15.49	0.198
Leucocytes/nl	8.21 ± 2.88	8.08 ± 2.50	0.743	8.42 ± 3.22	8.00 ± 2.48	0.155
Blood Met (%)	20.99 ± 8.07	22.31 ± 5.11	0.143	21.43 ± 7.36	21.27 ± 7.56	0.811
Met SAT (%)	11.95 ± 5.86	16.34 ± 6.54	0.011	13.56 ± 7.38	12.69 ± 5.83	0.784
Met VAT (%)	17.04 ± 5.61	19.69 ± 6.10	0.038	18.20 ± 4.41	17.46 ± 6.18	0.401
SAT <i>HIF3A</i> mRNA	21.08 ± 72.62	7.43 ± 40.53	0.209	11.58 ± 49.82	22.60 ± 76.47	0.660
VAT <i>HIF3A</i> mRNA	23.92 ± 106.19	10.45 ± 50.03	0.073	16.80 ± 82.25	24.09 ± 106.69	0.729

Table 3. Association of rs8102595 and rs3826795 with anthropometric and metabolic characteristics, mRNA expression and DNA methylation. Due to the low minor allele frequency (MAF) of the studied polymorphisms, subjects homozygous for the minor alleles (n = 3 for rs8102595, n = 16 for rs3826795) were combined with heterozygous groups (i.e. dominant mode of inheritance was used for statistical analyses). p-value adjusted for age, gender and BMI and diabetes status; BMI – Body Mass Index, WHR – waist-to-hip ratio, SAT – subcutaneous adipose tissue, VAT – visceral adipose tissue CRP – C-reactive protein, Met Blood (%) / Met SAT (%) / Met VAT (%) – Methylation of cg22891070 in *HIF3A* in blood/SAT/VAT, *HIF3A* mRNA – mRNA expression of *HIF3A* in subcutaneous/visceral adipose tissue.

high density lipoprotein (HDL)-cholesterol, gamma glutamyltransferase and the mRNA expression with PPARG ($p < 0.05$; Table 2).

After adjusting for age and gender, correlations between visceral and subcutaneous adipose tissue *HIF3A* mRNA expression and BMI, but also between visceral *HIF3A* mRNA expression and %body fat remained significant (Table 2). Correlations between subcutaneous and visceral adipose tissue *HIF3A* mRNA expression and waist, WHR, CRP level, leucocyte count and *leptin* mRNA expression remained significant after adjusting for age, gender and BMI (Table 2). After adjusting for covariates, free fatty acids only correlated with subcutaneous *HIF3A* mRNA expression and visceral *HIF3A* mRNA levels correlated with visceral mRNA expression of PPARG. In both fat depots, we found decreased *HIF3A* mRNA expression with increasing subcategories of both CRP serum concentrations and leucocyte counts (Fig. 2). To avoid a potential bias of systemic inflammation on *HIF3A* expression and its associations with anthropometric and metabolic traits, we performed correlation analyses only in individuals with CRP < 5mg/dl; however, the data remained unchanged (data not shown).

Association of rs8102595 and rs3826795 with *HIF3A* DNA methylation, mRNA expression and metabolic traits. In the present study, we included 2 SNPs (rs8102595 and rs3826795) which have previously been shown to be associated with DNA methylation in a large cohort including >2000¹¹. Both studied polymorphisms were in Hardy-Weinberg Equilibrium ($p > 0.05$) with following minor allele frequencies: rs8102595-10.8%, rs3826795-21.6%. There was no significant association between the SNPs and *HIF3A* mRNA expression in any of the two adipose tissue depots (Table 3). However, rs8102595 was nominally associated with *HIF3A* DNA methylation in visceral and subcutaneous adipose tissue ($p < 0.05$ after adjusting for age, gender and BMI; Table 3). Subjects carrying the minor allele (G) had a higher *HIF3A* DNA methylation in visceral adipose tissue, which was in line with the lower *HIF3A* mRNA expression in visceral adipose tissue (albeit not significant). Association analyses with parameters of obesity and fat distribution revealed a nominal association between rs3826795 and total cholesterol and the mean fat cell size of visceral adipose tissue (Supplementary Table 2). Rs8102595 showed an association with HDL-cholesterol, glucose infiltration rate and maximum fat cell size of subcutaneous adipose tissue (Table 3 and Supplementary Table 2).




Figure 3. Methylation of cg22891070 in *HIF3A* in different tissues and in relation to BMI (n = 87). The *HIF3A* locus is significantly higher methylated in blood compared to SAT and VAT (A) The difference in methylation levels can be observed for all BMI groups (B) Methylation levels are higher in subjects with increased BMI (B). Data are presented as means \pm SEM. * $p < 0.05$, ** $p < 0.01$, *** $p < 0.001$.

***HIF3A* DNA methylation in blood, subcutaneous and visceral adipose tissue.** Methylation measured at the CpG site in Assay 2, corresponds to the published cg22891070, which has been reported to show the strongest correlation to BMI¹¹. In our study, *HIF3A* DNA methylation at cg22891070 was significantly higher in blood ($20.84 \pm 7.74\%$) compared to subcutaneous ($12.83 \pm 6.82\%$; $p < 0.001$) and visceral adipose tissue ($17.28 \pm 5.61\%$; $p < 0.001$), whereas *HIF3A* DNA methylation in visceral adipose tissue was significantly higher than in subcutaneous adipose tissue ($p < 0.001$; Fig. 3A). In addition, DNA methylation at cg22891070 in visceral adipose tissue correlated significantly with hip ($p < 0.01$, $r = 0.614$), subcutaneous ($p < 0.01$, $r = 0.651$) and visceral fat mass ($p < 0.05$, $r = 0.468$) and inversely with the CT-ratio ($p < 0.01$, $r = -0.653$). Correlations between methylation in visceral adipose tissue and subcutaneous fat mass ($p < 0.01$), CT ratio ($p < 0.01$), hip ($p < 0.01$) and adiponectin ($p < 0.05$, $r = -0.187$) remained significant even after adjusting for age, gender and BMI. Furthermore, methylation of cg22891070 in subcutaneous adipose tissue correlated with CT ratio ($p < 0.05$, $r = -0.571$) and age ($p < 0.05$, $r = -0.268$). After adjusting for gender and BMI the correlation remained significantly for age ($p = 0.032$). Albeit not significant, in all analyzed tissues, obese individuals displayed a higher methylation of cg22891070 compared to lean and overweight individuals (Fig. 3B). The analyses including other tested CpG sites did not reveal correlations beyond those observed for cg22891070 (data not shown).

Discussion

Recent studies revealed an association between BMI and methylation of *HIF3A* in whole blood and in adipose tissue^{11–13}. It has been proposed that the HIF-system could play a role in mechanisms involved in the pathophysiology of adipose tissue-inflammation, obesity-induced insulin resistance and the etiology of obesity related diseases. We therefore sought to further elucidate the relationship between *HIF3A* mRNA expression in visceral and subcutaneous adipose tissue and obesity, but also methylation of CpG-sites in *HIF3A*. In summary, we show that *HIF3A* gene expression and methylation in adipose tissue are fat depot specific, and related to obesity and adipose tissue dysfunction.

We investigated the methylation and expression of *HIF3A* in two distinct fat depots, subcutaneous and visceral adipose tissue. We show that higher *HIF3A* mRNA expression in both subcutaneous and visceral adipose tissue is associated with higher BMI and obesity related traits. *HIF3A* has been shown to accelerate 3T3-L1 adipocyte differentiation and to induce the expression of adipocyte related genes⁹. Interestingly, we found higher adipose tissue *HIF3A* mRNA expression in individuals of the highest decile of mean adipocyte size (for both depots) compared to the lowest decile. This may suggest that *HIF3A* is involved in the determination of adipocyte size and may thereby contribute to adipose tissue expandability. Our results further support the hypothesis that expression of *HIF3A* might be induced in states of metabolic excess and mediate mechanisms involved in adipogenesis. Moreover, based on our data, the expression of *HIF3A* seems to be more pronounced in adipocytes compared to the stromal vascular fraction independent of the fat depot. To this end, adipocytes isolated from subcutaneous adipose tissue displayed higher expression levels of *HIF3A* than those isolated from visceral adipose tissue. Thus, the major proportion of *HIF3A* expression in adipose tissue might be attributed to primary adipocytes, which further supports the proposed regulatory role of *HIF3A* in adipogenesis. In further support of this, we found a positive correlation between the mRNA expression of *HIF3A* and *leptin* (in both visceral and subcutaneous adipose tissue) as well as *PPARG* (in visceral adipose tissue), two genes involved in the regulation of adipogenesis.

It is noteworthy, that *HIF3A* expression inversely correlated with CRP level and leucocyte count, suggesting down-regulation of the *HIF3A* expression in inflammatory states. Chronic inflammation in adipose tissue, liver and skeletal muscle are commonly associated with obesity², which results in secondary pathologies like insulin resistance, hyperinsulinemia and glucose intolerance^{3,21}. Obesity promoted relative hypoxia in adipocytes stimulates HIF1A-induction^{5,6}, which then triggers the inflammation process by mediating the production of adipocyte-derived chemokines and adipose tissue macrophage accumulation^{4,7}. *HIF3A* can inhibit HIF1A mediated signaling under certain circumstances²². The observed reduced expression of *HIF3A* in inflammatory states may facilitate increased HIF1A signaling, which in turn could activate an inflammatory cascade within adipose tissue.

HIF3A mRNA expression is regulated at different levels. Transcription of *HIF3A* can be induced by HIF1 via hypoxia response elements (HREs) in the promoter region and protein stability of HIF3 α can be regulated in dependency of oxygen supply via the oxygen-dependent degradation domain (ODD)^{22–24}. *HIF3A* expression has further been shown to be regulated by micro RNA (miRNA), thus to be modified on a post-transcriptional level²⁵. These different mechanisms can supplement one another in fine tuning of *HIF3A* expression. We hypothesize that the complex regulation of *HIF3A* expression can be influenced by DNA methylation in various ways by interfering with different mechanisms of regulation. An association between BMI and methylation at three CpG-sites in intron 1 of *HIF3A* in whole blood and in adipose tissue has recently been identified by employing genome-wide DNA-methylation analyses^{11–13}. In contrast, we did not find a correlation between BMI and *HIF3A* methylation. This may be due to the smaller sample size and a different composition of our cohort, which is characterized by a relatively high BMI (32.9 kg/m²), and thus, strongly differing from the previously reported cohorts with average BMI ranging between 24.2 and 28.3 kg/m². Rönn *et al.* were able to replicate the association between methylation of *HIF3A* and BMI in a female cohort only¹² and Demerath *et al.* showed *HIF3A* methylation to be associated with BMI only in one of three cohorts investigated¹³. Considering multiple isoforms of HIF3 α ⁸, it is plausible that methylation might be transcript-specific; yet, one would expect to observe consistent results upon expression analysis of the same transcript.

It is of note that the CpG sites at the *HIF3A* locus that were associated with BMI are situated within regions of open chromatin, suggesting that these sites lie in a regulatory region^{11,26}. However, this regulation appears more complex than being dependent on methylation only. It is plausible that methylation of *HIF3A* results in altered expression profiles, networking with mechanisms in different stages of regulation. Yet, a linear effect between methylation and expression even of the same transcript cannot be confirmed.

Methylation analysis of *HIF3A* in our cohort revealed significant differences between methylation in blood, subcutaneous and visceral adipose tissue, being strongest in blood and weakest in subcutaneous adipose tissue. Since *HIF3A* mRNA expression in subcutaneous adipose tissue is higher than in visceral adipose tissue, it is possible that methylation could together with other regulatory mechanisms, cause a decrease in the expression of *HIF3A*. In line with this, rs8102595 was nominally associated with DNA methylation at cg22891070 in subcutaneous and visceral adipose tissue; thus supporting data by Dick *et al.*¹¹ reporting associations of 2 SNPs (rs8102595 and rs3826795) with DNA methylation. Based on the Regulome Database²⁰, rs3826795 might affect the binding of transcription factors POLR2A and SIN3A, and rs8102595 might influence DNA-protein binding. However, considering the lack of associations of the two SNPs with BMI, changes in *HIF3A* methylation seem to be mediated by obesity rather than promoting obesity itself¹¹. It is also of note, that we did not observe an association between the SNPs and *HIF3A* mRNA expression in any of the two adipose tissue depots. We have to point out however, that the availability of the biomaterial (adipose tissue and blood samples) only allowed including 548 subjects for genotyping and for subsequent genotype-expression association analyses, which may have resulted in the lack of statistical power for correlation analyses.

In contrast to previous studies mostly investigating subcutaneous adipose tissue, the present study reveals mRNA expression and DNA methylation differences between subcutaneous and visceral adipose tissue. The two depots consist of different histological and biochemical compounds. The depot-specific expression of *HIF3A* may be important for the different functioning of the different depots. Whereas visceral adipose tissue is more vascular, innervated and contains a higher number of inflammatory and immune cells, subcutaneous adipose tissue has a higher preadipocyte differentiating capacity and a lower percentage of large adipocytes²⁷. As *HIF3A* mRNA expression is higher in subcutaneous adipose tissue, possibly due to differences in methylation, this contributes to our assumption that HIF3 α might be involved in preadipocyte differentiation, and that this process may be regulated by methylation, along with other factors. It is noteworthy that recently, we observed diminished hydroxymethylation levels in subcutaneous adipose tissue, as a measure of potential de-methylation mechanisms, which might be related to the higher number of pre-adipocytes in subcutaneous adipose tissue²⁸.

We found methylation of *HIF3A* in both compartments to be correlated inversely with fat distribution, and methylation in VAT correlated significantly with subcutaneous fat mass. This suggests that methylation occurs rather in subjects with a preponderance of subcutaneous fat. We also detected an inverse association between age and methylation in subcutaneous adipose tissue, which leads to the assumption that the modification is dynamic and changes during lifetime.

Finally, it has to be acknowledged that the CpG site cg22891070 presented in our study is located between the 2 previously reported CpG islands¹¹. Various *HIF3A* transcripts with different functions have been reported²² and it is also likely that they can be specifically affected by the methylation. Since the expression assay used in the present study tagged all potential *HIF3A* transcripts, we were not able to link cg22891070 to a specific transcript. However, in our own datasets based on genome-wide expression arrays (unpublished data) transcript variants 2 (NM_022462.4) and 3 (NM_152795.3) seem to be predominantly expressed in adipose tissue. Since Pasanen *et al.*²² suggested no functional relevance of the variant 3, it remains to be determined whether transcript variant 2 appears functionally relevant in adipose tissue.

In conclusion, our data suggest that *HIF3A* expression and methylation in adipose tissue is related to its dysfunction, making HIF3A an important factor involved in the complex etiology of obesity and associated comorbidities. HIF3A might function as an accelerator of adipogenesis in situations of excess of energetic supply and might contribute to the etiology of secondary obesity-induced pathologies by allowing a stronger induction of HIF1 α -mediated proinflammatory signaling.

References

1. Swinburn, B. A. *et al.* The global obesity pandemic: shaped by global drivers and local environments. *Lancet* **378**, 804–814 (2011).
2. Glass, C. K. & Olefsky, J. M. Inflammation and lipid signaling in the etiology of insulin resistance. *Cell Metab.* **15**, 635–645 (2012).
3. Shu, C. J., Benoist, C. & Mathis, D. The immune system's involvement in obesity-driven type 2 diabetes. *Semin. Immunol.* **24**, 436–442 (2012).

4. Lee, Y. S. *et al.* Increased adipocyte O₂ consumption triggers HIF-1 α , causing inflammation and insulin resistance in obesity. *Cell* **157**, 1339–1352 (2014).
5. Greer, S. N., Metcalf, J. L., Wang, Y. & Ohh, M. The updated biology of hypoxia-inducible factor. *EMBO J.* **31**, 2448–2460 (2012).
6. Majmundar, A. J., Wong, W. J. & Simon, M. C. Hypoxia-inducible factors and the response to hypoxic stress. *Mol. Cell* **40**, 294–309 (2010).
7. Gonsalves, C. S. & Kalra, V. K. Hypoxia-mediated expression of 5-lipoxygenase-activating protein involves HIF-1 α and NF- κ B and microRNAs 135a and 199a-5p. *J. Immunol.* **184**, 3878–3888 (2010).
8. Hara, S., Hamada, J., Kobayashi, C., Kondo, Y. & Imura, N. Expression and characterization of hypoxia-inducible factor (HIF)-3 α in human kidney: suppression of HIF-mediated gene expression by HIF-3 α . *Biochem. Biophys. Res. Commun.* **287**, 808–813 (2001).
9. Heidbreder, M. *et al.* Non-hypoxic induction of HIF-3 α by 2-deoxy-D-glucose and insulin. *Biochem. Biophys. Res. Commun.* **352**, 437–443 (2007).
10. Hatanaka, M. *et al.* Hypoxia-inducible factor-3 α functions as an accelerator of 3T3-L1 adipose differentiation. *Biological & pharmaceutical bulletin* **32**, 1166–1172 (2009).
11. Demerath, E. W. *et al.* Epigenome-wide association study (EWAS) of BMI, BMI change and waist circumference in African American adults identifies multiple replicated loci. *Human Molecular Genetics* **24**, 4464–4479 (2015).
12. Rönn, T. *et al.* Impact of age, BMI and HbA1c levels on the genome-wide DNA methylation and mRNA expression patterns in human adipose tissue and identification of epigenetic biomarkers in blood. *Human Molecular Genetics* **24**, 3792–3813 (2015).
13. Dick, K. J. *et al.* DNA methylation and body-mass index: a genome-wide analysis. *Lancet* **383**, 1990–1998 (2014).
14. Pan, H. *et al.* HIF3A association with adiposity: the story begins before birth. *Epigenomics* **7**, 1–13 (2015).
15. Huang, T. *et al.* DNA methylation variants at HIF3A locus, B vitamins intake, and long-term weight change: gene-diet interactions in two US cohorts. *Diabetes* **64**, 3146–3154 (2015).
16. American Diabetes Association. Diagnosis and classification of diabetes mellitus. *Diabetes care* **29** Suppl 1, S43–8 (2006).
17. Blüher, M. *et al.* Fas and FasL expression in human adipose tissue is related to obesity, insulin resistance, and type 2 diabetes. *The Journal of clinical endocrinology and metabolism* **99**, E36–44 (2014).
18. Klötting, N. *et al.* Insulin-sensitive obesity. *American journal of physiology. Endocrinology and metabolism* **299**, E506–15 (2010).
19. Blüher, M. *et al.* Adipose tissue selective insulin receptor knockout protects against obesity and obesity-related glucose intolerance. *Developmental cell* **3**, 25–38 (2002).
20. Boyle, A. P. *et al.* Annotation of functional variation in personal genomes using RegulomeDB. *Genome research* **22**, 1790–1797 (2012).
21. Lee, Y. S. *et al.* Inflammation is necessary for long-term but not short-term high-fat diet-induced insulin resistance. *Diabetes* **60**, 2474–2483 (2011).
22. Pasanen, A. *et al.* Hypoxia-inducible factor (HIF)-3 α is subject to extensive alternative splicing in human tissues and cancer cells and is regulated by HIF-1 but not HIF-2. *The international journal of biochemistry & cell biology* **42**, 1189–1200 (2010).
23. Tanaka, T., Wiesener, M., Bernhardt, W., Eckardt, K.-U. & Warnecke, C. The human HIF (hypoxia-inducible factor)-3 α gene is a HIF-1 target gene and may modulate hypoxic gene induction. *Biochem. J.* **424**, 143–151 (2009).
24. Heikkilä, M., Pasanen, A., Kivirikko, K. I. & Myllyharju, J. Roles of the human hypoxia-inducible factor (HIF)-3 α variants in the hypoxia response. *Cellular and molecular life sciences: CMLS* **68**, 3885–3901 (2011).
25. Gits, C. M. M. *et al.* MicroRNA response to hypoxic stress in soft tissue sarcoma cells: microRNA mediated regulation of HIF3 α . *BMC cancer* **14**, 429 (2014).
26. Cockerill, P. N. Structure and function of active chromatin and DNase I hypersensitive sites. *The FEBS journal* **278**, 2182–2210 (2011).
27. Ibrahim, M. M. Subcutaneous and visceral adipose tissue: structural and functional differences. *Obesity reviews: an official journal of the International Association for the Study of Obesity* **11**, 11–18 (2010).
28. Rohde, K. *et al.* DNA 5-hydroxymethylation in human adipose tissue differs between subcutaneous and visceral adipose tissue depots. *Epigenomics* **7**, 911–920 (2015).

Acknowledgements

We thank all those who participated in the study. This work was supported by grants from the IFB AdiposityDiseases (ADI-K50D and ADI-K7-45 to Y.B. and AD2-060E to P.K.) funded by the Federal Ministry of Education and Research (BMBF), Germany, FKZ: 01EO1501. This project was further supported by grants from the Collaborative Research Center funded by the German Research Foundation (CRC 1052; B01, B03, B04 to M.B., P.K. and N.K. respectively) and individual grant (HA 1374/15-1 to T.H.), by the Kompetenznetz Adipositas (Competence network for Obesity) funded by the German Federal Ministry of Education and Research (German Obesity Biomaterial Bank; FKZ 01GI1128) and by the German Diabetes Foundation (Deutsche Diabetes-Stiftung). Dorit Schleinitz is funded by the Boehringer Ingelheim Foundation.

Author Contributions

S.P., J.K., A.M. and N.E.H. performed mRNA expression and DNA methylation experiments; Y.B., N.K., T.H. and P.K. designed the study; M.R.S., A.D., M.F., T.L., M.D. and M.B. collected and provided biomaterial and clinical phenotypes; S.P., J.K., D.S., M.S., M.B. and P.K. wrote the manuscript.

Additional Information

Supplementary information accompanies this paper at <http://www.nature.com/srep>

Competing financial interests: The authors declare no competing financial interests.

How to cite this article: Pfeiffer, S. *et al.* Hypoxia-inducible factor 3A gene expression and methylation in adipose tissue is related to adipose tissue dysfunction. *Sci. Rep.* **6**, 27969; doi: 10.1038/srep27969 (2016).



This work is licensed under a Creative Commons Attribution 4.0 International License. The images or other third party material in this article are included in the article's Creative Commons license, unless indicated otherwise in the credit line; if the material is not included under the Creative Commons license, users will need to obtain permission from the license holder to reproduce the material. To view a copy of this license, visit <http://creativecommons.org/licenses/by/4.0/>

3 Summary

Dissertation in order to obtain the academic degree Dr. med.

***Hypoxia-inducible factor 3A* gene expression and methylation in adipose tissue is related to adipose tissue dysfunction**

Submitted by: Susanne Erna Margarete Pfeiffer

Submitted to the Faculty of Medicine of the University of Leipzig

Supervising tutor: Prof. Dr. med. Matthias Blüher

Date: 25.08.2016

Introduction

Obesity and its associated comorbidities constitute an evolving health burden worldwide (113). Obesity conditions chronic inflammation in adipose tissue, liver and skeletal muscle (44), which may contribute to chronic systemic inflammation, insulin resistance, and deterioration in glucose and lipid metabolism (46).

Upon weight gain, adipocyte hypertrophy may lead to hypoxia in adipose tissue (AT) which could be considered as a causative factor in the development of AT inflammation (45, 33, 114, 43). In this context, it has been recently shown that AT expression of *hypoxia inducible factor 1A* (*HIF1A*), a hypoxia inducible transcription factor, increases in mice early upon weight gain (52). In states of relative adipocyte hypoxia, induction of *HIF1A* (53, 54), stimulates accumulation of AT macrophages (55, 52) and the production of adipocyte-derived pro-inflammatory cytokines.

Hypoxia-inducible factors are transcription factors that mediate hypoxia response in various tissues (53). They are heterodimeric transcription factors consisting of an oxygen-labile α -subunit and a constitutively expressed β -subunit. Three different isoforms of the α -subunit, *HIF1 α* , *HIF2 α* and *HIF3 α* , exist and allow the formation of transcription factors with

different functions upon dimerizing with HIF β . The HIF1 α and HIF2 α target genes allow regulation of various biological processes such as erythropoiesis, angiogenesis, metabolic reprogramming, cell-cycle-regulation, and tumor progression, displaying master regulators of the transcriptional response to hypoxia (56, 57). HIF3 α structurally and functionally differs from HIF1 α /2 α (71, 72). According to a recent study, HIF3 α is capable of activating certain target genes independent or in collaboration with HIF1 α , suggesting a role of HIF3 α in glucose and amino acid metabolism, apoptosis, proteolysis, p53 signaling and PPAR signaling (78). In addition, HIF3 α has been shown to play a role in adipocyte differentiation (79).

Recent genome-wide analyses of DNA methylation in whole blood and human adipose tissue revealed an association of methylation at three CpG sites in intron 1 of *HIF3A* with BMI (104–106). In addition, two single nucleotide polymorphisms (SNPs) have been shown to be associated with methylation at these sites, yet to be independent of BMI (105). Further investigation of the relationship between hypoxia inducible factors and development of obesity-associated comorbidities might reveal important insights in pathophysiological processes concerning adipose tissue inflammation and/or insulin resistance in the etiology of obesity related metabolic diseases.

Objective

We therefore tested the hypothesis that expression of *HIF3A* in human subcutaneous and visceral AT is related to BMI, parameters of fat distribution and AT function, metabolic traits, genetic variation and methylation of CpG-sites in *HIF3A*.

Design and Methods

In paired samples of subcutaneous AT (SAT) and visceral AT (VAT) from 603 individuals with a wide range of age, BMI and obesity-related parameters, we investigated whether *HIF3A* mRNA expression is fat depot-specific, altered in obesity and related to measures of AT function and insulin sensitivity. We further tested for correlations between mean adipocyte size and AT *HIF3A* mRNA expression. In a subgroup of 548 individuals, we investigated the effects of *HIF3A* genetic variants on *HIF3A* AT expression and *HIF3A* methylation of CpG-sites. Further, we analyzed *HIF3A* mRNA expression in isolated adipocytes and cells from the stromal vascular fraction (SVF, i.e. preadipocytes, endothelial cells, endothelial smooth muscle cells, fibroblasts, macrophages, and blood cells isolated from adipose tissue). Methods are described in detail in the supplements.

Results

HIF3A expression is significantly higher in SAT compared to VAT and correlates in both fat depots with parameters of obesity, glucose metabolism and mRNA expression of genes regulating adipogenesis. AT *HIF3A* expression negatively correlates with circulating CRP and leucocytes independently of BMI ($p < 0.001$). *HIF3A* methylation at cg22891070 is significantly higher in VAT compared to SAT and correlates with BMI, abdominal SAT and VAT area. Rs8102595 shows a significant association with subcutaneous and visceral *HIF3A* methylation levels and rs3826795 and rs8102595 associate with obesity and fat distribution related traits. *HIF3a* mRNA expression in individuals with the highest decile of mean adipocyte size is higher compared to the lowest decile in both depots. We found significantly higher *HIF3A* expression in adipocytes compared to cells of the SVF in both SAT and VAT.

Discussion

We show that *HIF3A* gene expression and methylation in adipose tissue are fat depot specific, related to obesity and adipose tissue dysfunction parameters. Higher *HIF3A* mRNA expression in both subcutaneous and visceral adipose tissue is associated with higher BMI and obesity related traits. Our results further support the hypothesis that *HIF3A* signaling might be involved in adipogenesis and in the determination of adipocyte size. *HIF3A* expression is more pronounced in adipocytes compared to cells of the stromal vascular fraction, suggesting that the major proportion of *HIF3A* expression in adipose tissue might be attributed to primary adipocytes. We were also able to show a positive correlation between the mRNA expression of *HIF3A* and *leptin* (in both visceral and subcutaneous adipose tissue) and *PPARG* (in visceral adipose tissue), further supporting the proposed regulatory role of HIF3 α in adipogenesis. Our data support the hypothesis that hypoxia inducible transcription factor pathways may play an important role in the development of AT dysfunction associated with obesity. *HIF3A* gene expression inversely correlated with CRP level and leucocyte count, suggesting a down-regulation of *HIF3A* expression in inflammatory states. As HIF3 α can inhibit HIF1 α mediated signaling under certain circumstances, the reduced expression of *HIF3A* in inflammatory states may facilitate increased *HIF1A* signaling, which in turn activates the inflammatory cascade within adipose tissue.

HIF3A mRNA expression is regulated at different levels. Methylation analysis of *HIF3A* in our cohort revealed significant differences between methylation in blood, subcutaneous and

visceral adipose tissue, being strongest in blood and weakest in subcutaneous adipose tissue. Since *HIF3A* mRNA expression in subcutaneous adipose tissue is higher in visceral adipose tissue, it is possible that methylation could together with other regulatory mechanisms, cause a decrease in the expression of *HIF3A*. We found methylation of *HIF3A* in both compartments to be correlated inversely with fat distribution, and methylation in VAT correlated significantly with subcutaneous fat mass. This suggests that methylation occurs rather in subjects with a preponderance of subcutaneous fat. We also detected an inverse association between age and methylation in subcutaneous adipose tissue, which leads to the assumption that the modification is dynamic and changes during lifetime.

rs8102595 was associated with DNA methylation at cg22891070 in subcutaneous and visceral adipose tissue. However, we did not observe an association between rs8102595 or rs3826795 with BMI, suggesting that changes in *HIF3A* methylation seem to be mediated by obesity rather than promoting obesity itself.

In conclusion, our data suggest that *HIF3A* expression and methylation in adipose tissue is related to its dysfunction, making HIF3 α an important factor in the complex etiology of obesity and associated comorbidities.

4 Supplemental material

4.1 Material and methods

Analysis of human *HIF3A* mRNA expression

Briefly, human *HIF3A* mRNA expression was measured by qRT-PCR using TaqMan Gene Expression Assay (Applied Biosystems, Darmstadt, Germany), and fluorescence was detected on a TaqMan Quant Studio 6 Flex Real-Time PCR-System (Applied Biosystems, Darmstadt, Germany). Total RNA was isolated from AT samples using the Qiacube System (Qiagen, Hilden, Germany), and 2 µg RNA were reverse transcribed with standard reagents (Life Technologies). From each RT-PCR, 2 µl was amplified in a 20 µl PCR using the Taqman Gene Expression Assay (Applied Biosystems, Darmstadt, Germany) and the TaqMan Fast Advanced Mastermix (Applied Biosystems, Darmstadt, Germany). Samples were incubated in the Quant Studio 6 Flex Real-Time PCR-System (Applied Biosystems, Darmstadt, Germany) for an initial denaturation at 95°C for 20s, followed by 45 PCR cycles, each cycle consisting of 95°C for 1s and 60°C for 20s. The following Gene Expression Assay was used: Hs00541709_M1 (tagging the transcripts NM_022462.4, NM_152794.3, NM_152795.3 and NM_152796.4). *HIF3A* mRNA expression was calculated relative to the mRNA expression of *HPRT1*, determined by a premixed assay on demand for *HPRT1* mRNA (Hs01003267_M1, Applied Biosystems, Darmstadt, Germany). Expression of *HIF3A* and *HPRT1* mRNA were quantified by using the second derivative maximum method of the TaqMan Software (Applied Biosystems).

For expression analysis of *HIF3A* in adipocytes and SVF, total RNA was isolated from adipocytes and SVF extracted from paired samples of SAT and VAT. 305 ng RNA were reverse transcribed with standard reagents and from each RT-PCR, 23.5 µl was amplified in a 40 µl PCR using the Taqman Gene Expression Assay and the TaqMan Fast Advanced Mastermix according to the manufacturer's instruction. *HIF3A* mRNA expression was calculated relative to the mRNA expression of hypoxanthine guanine phosphoribosyltransferase 1 (*HPRT1*) mRNA or *18S* rRNA (for isolated adipocytes only), determined by a premixed assay on demand (Hs01003267_m1; Applied Biosystems, Darmstadt, Germany).

DNA extraction and bisulfite conversion

Genomic DNA was extracted using the DNeasy Blood and Tissue Kit (Qiagen, Hilden, Germany) and bisulfite conversion was performed using the Epitect Bisulfite Kit (Qiagen, Hilden, Germany) according to the manufacturer's protocol. PCRs were carried out to amplify DNA fragments for pyrosequencing. The following primers were used (Metabion, Martiensried, Germany): Forward 5'-TGGTTGAAGGGTTATTTAGGG-3'; reverse carrying a biotin label at its 5'-end 5'-ACTCTATCCCACCCCTTTT-3'. The PCR reaction mixture for pyrosequencing consisted of 5 µl 10x PCR buffer with MgCl₂ (Roche Diagnostics, Mannheim, Germany), 1 µl (10 mM dNTPs) PCR Grade Nucleotide Mix (Roche Diagnostics), 2.5 µl (10 pmol/µl) of forward and reverse primer (Metabion, München-Planegg, Germany), 0.4 µl (5 U/µl) FastStart Taq DNA Polymerase (Roche Diagnostics), 2 µl of bisulfite-converted DNA and 36.6 µl PCR-grade water. Amplifications were performed with an initial denaturation step at 95°C for 5 min, 38 cycles of 95°C for 30 s, 60°C for 30 s, and 72°C for 45 s, and a final extension step at 72°C for 5 min.

4.2 Results

Supplementary Table 1. Correlation analyses between subcutaneous and visceral adipose tissue *HIF3A* mRNA expression and study parameters.

	<i>HIF3A</i> mRNA Expression in subcutaneous adipose tissue			<i>HIF3A</i> mRNA Expression in visceral adipose tissue		
	r	p-value	adj. p-value	r	p-value	adj. p-value
Age (years)	-0.23	4.61x10 ⁻⁵	0.032	-0.237	3.08x10 ⁻⁵	0.076
BMI (kg/m ²)	0.239	2.86x10 ⁻⁵	0.017^a	0.283	5.46x10 ⁻⁷	8.84x10^{-4a}
Body weight (kg)	0.235	5.56x10 ⁻⁵	0.467 ^a	0.263	5.45x10 ⁻⁶	0.280 ^a
Height (m)	0.044	0.458	0.467	0.001	0.983	0.538
Waist (cm)	0.472	8.41x10 ⁻⁹	0.010	0.515	1.89x10 ⁻¹⁰	0.048
Hip (cm)	0.387	2.13x10 ⁻⁵	0.425	0.442	6.73x10 ⁻⁷	0.628
WHR	0.172	0.067	0.018	0.139	0.135	0.033
Visceral fat area (cm ²)	0.391	3.19x10 ⁻⁵	0.636	0.442	1.71x10 ⁻⁶	0.479
SC fat area (cm ²)	0.392	2.99x10 ⁻⁵	0.240	0.465	4.06x10 ⁻⁷	0.604
CT ratio (sc/vis)	-0.259	7.04x10 ⁻³	0.165	-0.319	7.80x10 ⁻⁴	0.325
Body fat (%)	0.324	0.017	0.055 ^a	0.442	8.23x10 ⁻⁴	0.013
CRP (mg/dl)	-0.138	0.021	1.8x10⁻³	-0.153	0.010	3.19x10⁻⁴
FPG (mmol/l)	0.077	0.200	0.180	0.017	0.770	0.784
FPI (pmol/l)	0.130	0.181	0.161	0.250	8.21x10 ⁻³	0.634
Total Cholesterol (mmol/l)	-0.017	0.827	0.673	-0.010	0.891	0.586
HDL-C (mmol/l)	-0.138	0.141	0.913	-0.211	0.022	0.602
LDL-C	0.109	0.249	0.896	0.111	0.236	0.802
FFA (mmol/l)	0.442	1.99x10 ⁻⁵	8.97x10⁻³	0.401	1.43x10 ⁻⁴	0.063
TG (mmol/l)	0.218	3.40x10 ⁻³	0.085	0.196	7.63x10 ⁻³	0.262
Leptin (ng/ml)	0.413	2.14x10 ⁻⁵	0.866	0.461	1.78x10 ⁻⁶	0.918
Adiponectin (μg/ml)	-0.290	3.47x10 ⁻³	0.415	-0.0348	4.09x10 ⁻⁴	0.241
Albumin (g/L)	-0.328	0.072	0.159	-0.217	0.225	0.015
ALAT (μkat/l)	0.176	3.31x10 ⁻³	0.303	0.135	0.023	0.980
gGT (μkat/l)	-0.101	0.095	0.226	-0.148	0.014	0.040
TSH (mU/l)	0.091	0.153	0.850	0.128	0.043	0.471
Leucocytes/nl	-0.127	0.032	3.05x10⁻³	-0.133	0.024	1.13x10⁻³
Met Blood (%)	0.054	0.720	0.618	0.023	0.876	0.772
Met SAT (%)	-0.054	0.687	0.482	-0.088	0.498	0.345
Met VAT (%)	0.060	0.648	0.667	-0.045	0.729	0.757


Supplementary table 1. r - correlation coefficient (Pearson adj. - p-value adjusted to age, gender and BMI, ^a adjusted for gender and age; BMI – Body Mass Index, WHR – waist-to-hip ratio, CRP – C-reactive protein, FPI – fasting plasma insulin, HDL-C – high density lipoprotein cholesterol, FFA – free fatty acids, TG – triglycerides, ALAT – alanine aminotransferase, gGT - gamma-glutamyl transferase, TSH – thyroid-stimulation hormone, Met Blood (%) / Met SAT (%) / Met VAT (%) - methylation of cg22891070 in *HIF3A* in blood / SAT/ VAT

Supplementary Table 2. Association of rs8102595 and rs3826795 with anthropometric and metabolic characteristics, mRNA expression and DNA methylation.

	rs8102595			rs3826795		
	A/A	A/G + G/G	p-value	A/A+ A/G	G/G	p-value
N	446	95		208	336	
Men/Women	151/295	32/63		73/135	110/226	
Age	52.83±15.79	55.48±15.44	0.482	49.56±15.31	50.72±14.69	0.278
BMI (kg/m ²)	43.48±13.74	42.51±13.50	0.239	43.64±14.04	42.93±13.32	0.908
Body weight (kg)	126.86±42.81	124.57±40.14	0.680	128.42±45.54	124.60±41.15	0.769
Height (m)	1.69±0.09	1.69±0.9	0.628	1.69±0.09	1.69±0.09	0.763
Waist (cm)	124.26±29.98	121.84±30.09	0.798	124.46±30.43	122.85±29.87	0.935
Hip (cm)	130.53±28.99	128.59±28.38	0.851	129.54±28.38	130.08±29.56	0.676
WHR	0.95±0.13	0.96±0.16	0.316	0.96±0.16	0.94±0.12	0.921
VAT area (cm ²)	242.93±173.84	237.02±159.92	0.575	256.05±183.40	228.98±159.97	0.674
SAT area (cm ²)	1095.74±795.48	1129.73±819.78	0.536	1122.85±774.80	1094.46±817.64	0.902
VAT mean	123.00±20.82	122.08±20.60	0.999	119.69±25.71	124.66±17.25	0.014
SAT mean	127.37±19.89	127.51±17.42	0.486	126.50±19.04	127.99±19.84	0.334
VAT max	209.23±58.51	230.21±96.06	0.060	210.73±74.84	213.66±63.47	0.109
SAT max	214.28±70.88	249.22±110.69	1.23x10⁻³	224.71±80.22	217.94±79.94	0.987
CT ratio (vis/sc)	0.47±0.63	0.38±0.30	0.922	0.40±0.42	0.48±0.66	0.826
Body fat (%)	41.95±11.35	42.26±11.72	0.496	41.15±11.88	42.57±11.11	0.607
CRP (mg/dl)	12.04±15.09	11.20±16.05	0.935	13.09±15.67	11.34±15.49	0.198
IL-6 (pg/ml)	6.05±5.24	5.17±3.71	0.880	6.16±4.54	5.79±5.39	0.286
HbA1C (%)	6.07±1.11	5.82±0.86	0.443	6.00±1.10	6.02±1.05	0.766
oGTT2h (mmol/l)	6.99±2.41	7.49±3.97	0.663	7.57±3.40	6.74±2.22	0.064
FPG (mmol/l)	6.34±2.17	5.83±0.98	0.827	6.34±2.45	6.22±1.77	0.904
FPI (pmol/l)	131.78±141.75	107.18±98.96	0.297	124.28±118.40	127.54±145.26	0.729
GIR (μmol/kg/min)	75.66±33.87	65.78±35.86	0.042	73.75±33.57	73.58±35.24	0.567
Total cholesterol (mmol/l)	4.87±0.99	4.72±0.97	0.130	4.99±1.06	4.75±0.93	0.019
HDL-C (mmol/l)	1.24±0.41	1.22±0.46	0.027	1.25±0.40	1.23±0.42	0.418
LDL-C (mmol/l)	3.11±0.94	3.02±1.12	0.143	3.19±1.02	3.01±0.93	0.167
FFA (mmol/l)	0.58±0.41	0.54±0.39	0.525	0.57±0.43	0.57±0.39	0.076
TG (mmol/l)	1.65±0.89	1.49±0.66	0.654	1.69±0.96	1.58±0.78	0.444
Leptin (ng/ml)	37.12±22.14	40.64±23.96	0.313	35.43±21.13	39.15±23.34	0.655
Adiponectin (μg/ml)	7.93±4.97	6.59±3.36	0.091	7.17±4.67	8.11±4.75	0.095
Albumin (g/l)	40.95±9.62	40.59±9.43	0.648	39.37±10.74	41.84±8.40	0.086
ALAT (μkat/l)	0.65±0.50	0.58±0.49	0.360	0.65±0.53	0.62±0.47	0.767
ASAT (μkat/l)	0.71±3.18	0.52±0.31	0.382	0.89±0.65	0.54±0.28	0.742
gGT (μkat/l)	0.89±1.65	0.81±0.77	0.679	0.85±1.33	0.88±1.64	0.960
TSH (mU/l)	2.317±10.87	1.735±2.64	0.296	2.96±15.74	1.73±2.06	0.071
fT3 (pg/ml)	4.66±0.99	4.71±0.81	0.348	4.76±1.03	4.62±0.92	0.609
fT4 (pmol/l)	17.15±3.47	17.81±3.63	0.354	16.78±2.68	17.65±3.71	0.222
Blood Met (%)	20.99±8.07	22.31±5.11	0.143	21.43±7.36	21.27±7.56	0.811
Met SAT (%)	11.95±5.86	16.34±6.54	0.011	13.56±7.38	12.69±5.83	0.784
Met VAT (%)	17.04±5.61	19.69±6.10	0.038	18.20±4.41	17.46±6.18	0.401
SAT <i>HIF3A</i> mRNA	21.08±72.62	7.43±40.53	0.209	11.58±49.82	22.60±76.47	0.660
VAT <i>HIF3A</i> mRNA	23.92±106.19	10.45±50.03	0.073	16.80±82.25	24.09±106.69	0.729
Leucocytes/nl	8.21±2.88	8.08±2.50	0.743	8.42±3.22	8.00±2.48	0.155
Erythrocytes (Mio/μl)	4.79±1.05	4.62±0.50	0.230	4.65±0.59	4.82±1.16	0.237
Thrombocytes x10 ⁹ /l	257.94±82.94	291.52±101.69	0.089	261.59±75.91	263.98±93.83	0.762

Supplementary table 2. Due to the low minor allele frequency (MAF) of the studied polymorphisms, subjects homozygous for the minor alleles (n=3 for rs8102595, n=16 for rs3826795) were combined with heterozygous groups (i.e. dominant mode of inheritance was used for statistical analyses). p-value adjusted for age, gender and BMI and diabetes status; BMI – Body Mass Index, WHR – waist-to-hip ratio, SAT – subcutaneous adipose tissue, CRP – C-reactive protein, IL-6 – Interleukin 6, HbA1c – glycohemoglobin, oGTT – oral glucose tolerance test, FPG – fasting plasma glucose, FPI – fasting plasma insulin, GIR – glucose infusion rate during the steady state of an euglycemic hyperinsulinemic clamp, HDL-C – high density lipoprotein cholesterol, LDL-C – low density lipoprotein cholesterol, FFA – free fatty acids, TG – triglycerides, ALAT – alanine aminotransferase, ASAT – aspartate aminotransferase, gGT – gamma-glutamyl transferase, TSH – thyroid-stimulation hormone, fT3 – free triiodothyronine, fT4 – free tetraiodothyronine, Met Blood (%) / Met Sc (%) / Met Visc (%) – methylation of cg22891070 in *HIF3A* in blood / SAT / VAT, *HIF3A* mRNA – mRNA expression of *HIF3A* in subcutaneous/visceral adipose tissue.

Supplementary Figure. Fat depot-related *HIF3A* mRNA expression pattern is distinct in individuals with either normal glucose tolerance (NGT; SAT, n=316; VAT, n=242) or type 2 diabetes (T2D; SAT, n=318; VAT, n=245)



Supplementary Figure. NGT- normal glucose tolerance; SAT – subcutaneous adipose tissue; VAT – visceral adipose tissue, T2D – type 2 diabetes

5 Declaration of the independent formulation of the work

Hiermit erkläre ich, dass ich die vorliegende Arbeit selbständig und ohne unzulässige Hilfe oder Benutzung anderer als der angegebenen Hilfsmittel angefertigt habe. Ich versichere, dass Dritte von mir weder unmittelbar noch mittelbar geldwerte Leistungen für Arbeiten erhalten haben, die im Zusammenhang mit dem Inhalt der vorgelegten Dissertation stehen, und dass die vorgelegte Arbeit weder im Inland noch im Ausland in gleicher oder ähnlicher Form einer anderen Prüfungsbehörde zum Zweck einer Promotion oder eines anderen Prüfungsverfahrens vorgelegt wurde. Alles aus anderen Quellen und von anderen Personen übernommene Material, das in der Arbeit verwendet wurde oder auf das direkt Bezug genommen wird, wurde als solches kenntlich gemacht. Insbesondere wurden alle Personen genannt, die direkt an der Entstehung der vorliegenden Arbeit beteiligt waren. Ich versichere, dass die aktuellen gesetzlichen Vorgaben in Bezug auf die Zulassung der klinischen Studien, die Bestimmungen des Tierschutzgesetzes, die Bestimmungen des Gentechnikgesetzes und die allgemeinen Datenschutzbedingungen eingehalten wurden und ein positives Ethikvotum vorliegt. Weiterhin versichere ich, dass die Regelungen zur Sicherung guter wissenschaftlicher Praxis eingehalten wurden.

25.08.2016

Susanne Pfeiffer

6 Curriculum vitae

Personal Data

Name	Susanne Erna Margarete Pfeiffer
Date of birth	02/04/1989
Place of birth	Munich, Germany
Marital Status	single

School Education

08/2000 - 06/2007	Werner-von-Siemens secondary school in Berlin
01/2005 - 06/2005	semester abroad at the Highschool of Stevensville, Michigan, USA
06/2007	successful accomplishment of the "Allgemeine Hochschulreife"

Medical Education

10/2007 - 06/2009	beginning of the academic studies of humane medicine at the Albert Szent-Györgyi University in Szeged, Hungary
10/2009 - 11/2013	continuation of the academic studies of humane medicine at the University of Leipzig
08/2012 – 12/2012	completion of the 1 st part of the „Praktisches Jahr“ (surgery) at the Hospital Universitario de la Princesa, Madrid, Spain
12/2012 – 04/2013	completion of the 2 nd and 3 rd part of the “Praktisches Jahr” (internal medicine und urology) at University Hospital of Leipzig
11/2013	successful accomplishment of the „Zweiter Abschnitt der Ärztlichen Prüfung“

Work Experience

09/2015 – 04/2016	Charité Berlin, Department of Surgery residency in surgery
from 05/2016	Charité Berlin, Department of Pediatrics, Clinic for pediatric endocrinology residency in pediatrics

Dissertation and Scientific Activity

- 07/2014 – 08/2015 dissertation at the IFB Adiposity and Disease of the University of Leipzig with Professor Dr. med. Matthias Blüher
- 12/2014- 08/2015 scientific project at the Helmholtz Center for environmental Research Leipzig with Prof. Dr. Martin von Bergen

Scientific Publications

- 06/2016 “Hypoxia-inducible factor 3A gene expression and methylation in adipose tissue is related to adipose tissue dysfunction” published in “Scientific Reports 2016; 6:27969” and written by “Pfeiffer S*, Krüger J*, Maierhofer A, Böttcher Y, Klötting N, El Hajj N, Schleinitz D, Schön MR, Dietrich A, Fasshauer M, Lohmann T, Dreßler M, Stummvoll M, Haaf T, Blüher M, Kovacs P (*These authors contributed equally to the work)
- 07/2016 “Microdialysis Sampling from Wound Fluids Enables Quantitative Assessment of Cytokines, Proteins, and Metabolites Reveals Bone Defect-Specific Molecular Profiles” published in “Plos One. 2016; 11:e0159580” and written by “Förster Y, Schmidt JR, Wissenbach DK, Pfeiffer Susanne EM, Baumann S, Hofbauer LC, Bergen M von, Kalkhof S, Rammelt S”

7 Acknowledgement

In erster Linie möchte ich meinem Doktorvater Prof. Dr. med. Matthias Blüher für die Möglichkeit der Anfertigung dieser interessanten Arbeit unter seiner Betreuung danken. Seine vielen klugen Ideen und Bereitschaft, das Thema vielseitig mit mir zu diskutieren und auf meine Fragen einzugehen, haben mich stets geleitet und motiviert. Mein Dank gilt außerdem Prof. Dr. Peter Kovacs, der mit großer Offenheit und Diskussionsbereitschaft die Anfertigung dieser Arbeit unterstützt hat.

Außerdem möchte ich besonders Jacqueline Krüger danken. Ihre Anleitungen, Hilfe und Geduld im Labor und bei der Auswertung der Daten sind mir eine große Unterstützung gewesen.

Ich möchte mich in besonderem Maße sehr herzlich bei Nora Klötting-Blüher, Matthias und Daniela Kern, Viola Döbel, Lutz Feige, Gesine Flehming, Nico Hesselbarth, Nicole Lakowa sowie allen anderen Mitarbeitern der Arbeitsgruppe Blüher für ihre Hilfsbereitschaft sowie angenehme und entspannte Arbeitsatmosphäre bedanken. Ich werde diese Zeit immer in guter Erinnerung behalten und gerne an sie zurückdenken.

Zum Abschluss gilt der größte Dank meinen Freunden, meinem Freund und meiner Familie, die mich in jeder Stimmungslage bedingungslos unterstützt haben.

8 References

1. **Speakman JR, O'Rahilly S** Fat: an evolving issue, *Dis Model Mech.* 2012; 5:569–573
2. **Adler M, Schaffner F** Fatty liver hepatitis and cirrhosis in obese patients, *Am. J. Med.* 1979; 67:811–816
3. **Angulo P** Nonalcoholic fatty liver disease, *N. Engl. J. Med.* 2002; 346:1221–1231
4. **Hofbauer KG, Nicholson JR** Pharmacotherapy of obesity, *Exp. Clin. Endocrinol. Diabetes.* 2006; 114:475–484
5. **Jeong S, Nam H, Son M, Son E, Cho K** Interactive effect of obesity indexes on cognition, *Dement Geriatr Cogn Disord.* 2005; 19:91–96
6. **Kahn SE, Hull RL, Utzschneider KM** Mechanisms linking obesity to insulin resistance and type 2 diabetes, *Nature.* 2006; 444:840–846
7. **LeRoith D, Novosyadlyy R, Gallagher EJ, Lann D, Vijayakumar A, Yakar S** Obesity and type 2 diabetes are associated with an increased risk of developing cancer and a worse prognosis; epidemiological and mechanistic evidence, *Exp. Clin. Endocrinol. Diabetes.* 2008; 116 Suppl 1:S4–6
8. **Sainaghi PP, Castello L, Bergamasco L, Carnevale Schianca, G P, Bartoli E** Metabolic characteristics of glucose intolerance: the critical role of obesity, *Exp. Clin. Endocrinol. Diabetes.* 2008; 116:86–93
9. **Van Gaal, Luc F, Mertens IL, De Block, Christophe E** Mechanisms linking obesity with cardiovascular disease, *Nature.* 2006; 444:875–880
10. **Blüher M** Are there still healthy obese patients?, *Curr Opin Endocrinol Diabetes Obes.* 2012; 19:341–346
11. **Müller MJ, Lagerpusch M, Enderle J, Schautz B, Heller M, Bosy-Westphal A** Beyond the body mass index: tracking body composition in the pathogenesis of obesity and the metabolic syndrome, *Obes Rev.* 2012; 13 Suppl 2:6–13
12. **Reaven GM** Importance of identifying the overweight patient who will benefit the most by losing weight, *Ann. Intern. Med.* 2003; 138:420–423
13. **Thomas EL, Parkinson JR, Frost GS, Goldstone AP, Doré CJ, McCarthy JP, Collins AL, Fitzpatrick JA, Durighel G, Taylor-Robinson SD, Bell JD** The missing risk: MRI and MRS phenotyping of abdominal adiposity and ectopic fat, *Obesity (Silver Spring).* 2012; 20:76–87
14. **Blüher M** Adipose tissue dysfunction in obesity, *Exp. Clin. Endocrinol. Diabetes.* 2009; 117:241–250
15. **Ohlson LO, Larsson B, Svärdsudd K, Welin L, Eriksson H, Wilhelmsen L, Björntorp P, Tibblin G** The influence of body fat distribution on the incidence of diabetes mellitus. 13.5 years of follow-up of the participants in the study of men born in 1913, *Diabetes.* 1985; 34:1055–1058

16. **Abate N, Garg A** Heterogeneity in adipose tissue metabolism: causes, implications and management of regional adiposity, *Prog. Lipid Res.* 1995; 34:53–70
17. **Mauriège P, Després JP, Moorjani S, Prud'Homme D, Lamarche B, Bouchard C, Nadeau A, Tremblay A, Lupien PJ** Abdominal and femoral adipose tissue lipolysis and cardiovascular disease risk factors in men, *Eur. J. Clin. Invest.* 1993; 23:729–740
18. **Ibrahim MM** Subcutaneous and visceral adipose tissue: structural and functional differences, *Obes Rev.* 2010; 11:11–18
19. **Poznanski WJ, Waheed I, Van R** Human fat cell precursors. Morphologic and metabolic differentiation in culture, *Lab Invest.* 1973; 29:570–576
20. **Blüher M** Adipokines - removing road blocks to obesity and diabetes therapy, *Mol Metab.* 2014; 3:230–240
21. **Catalán V, Gómez-Ambrosi J, Rodríguez A, Salvador J, Frühbeck G** Adipokines in the treatment of diabetes mellitus and obesity, *Expert Opin Pharmacother.* 2009; 10:239–254
22. **Ahima RS, Flier JS** Leptin, *Annu Rev Physiol.* 2000; 62:413–437
23. **Tang QQ, Lane MD** Adipogenesis: from stem cell to adipocyte, *Annu. Rev. Biochem.* 2012; 81:715–736
24. **Park KW, Halperin DS, Tontonoz P** Before they were fat: adipocyte progenitors, *Cell Metab.* 2008; 8:454–457
25. **Medina-Gomez G, Gray SL, Yetukuri L, Shimomura K, Virtue S, Campbell M, Curtis RK, Jimenez-Linan M, Blount M, Yeo, Giles S H, Lopez M, Seppänen-Laakso T, Ashcroft FM, Oresic M, Vidal-Puig A** PPAR gamma 2 prevents lipotoxicity by controlling adipose tissue expandability and peripheral lipid metabolism, *PLoS Genet.* 2007; 3:e64
26. **Wajchenberg BL** Subcutaneous and visceral adipose tissue: their relation to the metabolic syndrome, *Endocr. Rev.* 2000; 21:697–738
27. **Heinrich PC, Castell JV, Andus T** Interleukin-6 and the acute phase response, *Biochem. J.* 1990; 265:621–636
28. **Mårin P, Andersson B, Ottosson M, Olbe L, Chowdhury B, Kvist H, Holm G, Sjöström L, Björntorp P** The morphology and metabolism of intraabdominal adipose tissue in men, *Metab. Clin. Exp.* 1992; 41:1242–1248
29. **Enzi G, Gasparo M, Biondetti PR, Fiore D, Semisa M, Zurlo F** Subcutaneous and visceral fat distribution according to sex, age, and overweight, evaluated by computed tomography, *Am. J. Clin. Nutr.* 1986; 44:739–746
30. **Goss AM, Darnell BE, Brown MA, Oster RA, Gower BA** Longitudinal associations of the endocrine environment on fat partitioning in postmenopausal women, *Obesity (Silver Spring).* 2012; 20:939–944

31. **Bouchard C, Tremblay A, Després JP, Nadeau A, Lupien PJ, Thériault G, Dussault J, Moorjani S, Pinault S, Fournier G** The response to long-term overfeeding in identical twins, *N. Engl. J. Med.* 1990; 322:1477–1482
32. **Smith SR, Zachwieja JJ** Visceral adipose tissue: a critical review of intervention strategies, *Int. J. Obes. Relat. Metab. Disord.* 1999; 23:329–335
33. **Trayhurn P** Hypoxia and adipose tissue function and dysfunction in obesity, *Physiol. Rev.* 2013; 93:1–21
34. **Blaak EE, van Baak, M A, Kemerink GJ, Pakbiers MT, Heidendal GA, Saris WH** Beta-adrenergic stimulation and abdominal subcutaneous fat blood flow in lean, obese, and reduced-obese subjects, *Metab. Clin. Exp.* 1995; 44:183–187
35. **Kabon B, Nagele A, Reddy D, Eagon C, Fleshman JW, Sessler DI, Kurz A** Obesity decreases perioperative tissue oxygenation, *Anesthesiology.* 2004; 100:274–280
36. **Goossens GH, Bizzarri A, Venteclef N, Essers Y, Cleutjens JP, Konings E, Jocken, Johan W E, Cajlakovic M, Ribitsch V, Clément K, Blaak EE** Increased adipose tissue oxygen tension in obese compared with lean men is accompanied by insulin resistance, impaired adipose tissue capillarization, and inflammation, *Circulation.* 2011; 124:67–76
37. **Karpe F, Fielding BA, Ilic V, Macdonald IA, Summers, Lucinda K M, Frayn KN** Impaired postprandial adipose tissue blood flow response is related to aspects of insulin sensitivity, *Diabetes.* 2002; 51:2467–2473
38. **Skurk T, Alberti-Huber C, Herder C, Hauner H** Relationship between adipocyte size and adipokine expression and secretion, *J. Clin. Endocrinol. Metab.* 2007; 92:1023–1033
39. **Brahimi-Horn MC, Pouysségur J** Oxygen, a source of life and stress, *FEBS Lett.* 2007; 581:3582–3591
40. **Pasarica M, Sereda OR, Redman LM, Albarado DC, Hymel DT, Roan LE, Rood JC, Burk DH, Smith SR** Reduced adipose tissue oxygenation in human obesity: evidence for rarefaction, macrophage chemotaxis, and inflammation without an angiogenic response, *Diabetes.* 2009; 58:718–725
41. **Spencer M, Unal R, Zhu B, Rasouli N, McGehee RE, Peterson CA, Kern PA** Adipose tissue extracellular matrix and vascular abnormalities in obesity and insulin resistance, *J. Clin. Endocrinol. Metab.* 2011; 96:E1990-8
42. **Rausch ME, Weisberg S, Vardhana P, Tortoriello DV** Obesity in C57BL/6J mice is characterized by adipose tissue hypoxia and cytotoxic T-cell infiltration, *Int J Obes (Lond).* 2008; 32:451–463
43. **Ye J, Gao Z, Yin J, He Q** Hypoxia is a potential risk factor for chronic inflammation and adiponectin reduction in adipose tissue of ob/ob and dietary obese mice, *Am. J. Physiol. Endocrinol. Metab.* 2007; 293:E1118-28

44. **Glass CK, Olefsky JM** Inflammation and lipid signaling in the etiology of insulin resistance, *Cell Metab.* 2012; 15:635–645
45. **Klötting N, Blüher M** Adipocyte dysfunction, inflammation and metabolic syndrome, *Rev Endocr Metab Disord.* 2014; 15:277–287
46. **Shu CJ, Benoist C, Mathis D** The immune system's involvement in obesity-driven type 2 diabetes, *Semin. Immunol.* 2012; 24:436–442
47. **Hotamisligil GS, Shargill NS, Spiegelman BM** Adipose expression of tumor necrosis factor- α : direct role in obesity-linked insulin resistance, *Science.* 1993; 259:87–91
48. **Weisberg SP, McCann D, Desai M, Rosenbaum M, Leibel RL, Ferrante AW** Obesity is associated with macrophage accumulation in adipose tissue, *J. Clin. Invest.* 2003; 112:1796–1808
49. **Xu H, Barnes GT, Yang Q, Tan G, Yang D, Chou CJ, Sole J, Nichols A, Ross JS, Tartaglia LA, Chen H** Chronic inflammation in fat plays a crucial role in the development of obesity-related insulin resistance, *J. Clin. Invest.* 2003; 112:1821–1830
50. **Fève B, Bastard J** The role of interleukins in insulin resistance and type 2 diabetes mellitus, *Nat Rev Endocrinol.* 2009; 5:305–311
51. **Hotamisligil GS** Mechanisms of TNF- α -induced insulin resistance, *Exp. Clin. Endocrinol. Diabetes.* 1999; 107:119–125
52. **Lee YS, Kim J, Osborne O, Oh DY, Sasik R, Schenk S, Chen A, Chung H, Murphy A, Watkins SM, Quehenberger O, Johnson RS, Olefsky JM** Increased adipocyte O₂ consumption triggers HIF-1 α , causing inflammation and insulin resistance in obesity, *Cell.* 2014; 157:1339–1352
53. **Greer SN, Metcalf JL, Wang Y, Ohh M** The updated biology of hypoxia-inducible factor, *EMBO J.* 2012; 31:2448–2460
54. **Majmundar AJ, Wong WJ, Simon MC** Hypoxia-inducible factors and the response to hypoxic stress, *Mol. Cell.* 2010; 40:294–309
55. **Gonsalves CS, Kalra VK** Hypoxia-mediated expression of 5-lipoxygenase-activating protein involves HIF-1 α and NF- κ B and microRNAs 135a and 199a-5p, *J. Immunol.* 2010; 184:3878–3888
56. **Gordan JD, Simon MC** Hypoxia-inducible factors: central regulators of the tumor phenotype, *Curr. Opin. Genet. Dev.* 2007; 17:71–77
57. **Rankin EB, Giaccia AJ** The role of hypoxia-inducible factors in tumorigenesis, *Cell Death Differ.* 2008; 15:678–685
58. **Patel SA, Simon MC** Biology of hypoxia-inducible factor-2 α in development and disease, *Cell Death Differ.* 2008; 15:628–634

-
59. **Pugh CW, O'Rourke JF, Nagao M, Gleadle JM, Ratcliffe PJ** Activation of hypoxia-inducible factor-1; definition of regulatory domains within the alpha subunit, *J. Biol. Chem.* 1997; 272:11205–11214
 60. **Huang LE, Pete EA, Schau M, Milligan J, Gu J** Leu-574 of HIF-1alpha is essential for the von Hippel-Lindau (VHL)-mediated degradation pathway, *J. Biol. Chem.* 2002; 277:41750–41755
 61. **Kageyama Y, Koshiji M, To KKW, Tian Y, Ratcliffe PJ, Huang LE** Leu-574 of human HIF-1alpha is a molecular determinant of prolyl hydroxylation, *FASEB J.* 2004; 18:1028–1030
 62. **Bruick RK, McKnight SL** A conserved family of prolyl-4-hydroxylases that modify HIF, *Science.* 2001; 294:1337–1340
 63. **Epstein AC, Gleadle JM, McNeill LA, Hewitson KS, O'Rourke J, Mole DR, Mukherji M, Metzen E, Wilson MI, Dhanda A, Tian YM, Masson N, Hamilton DL, Jaakkola P, Barstead R, Hodgkin J, Maxwell PH, Pugh CW, Schofield CJ, Ratcliffe PJ** C. elegans EGL-9 and mammalian homologs define a family of dioxygenases that regulate HIF by prolyl hydroxylation, *Cell.* 2001; 107:43–54
 64. **Hirsilä M, Koivunen P, Günzler V, Kivirikko KI, Myllyharju J** Characterization of the human prolyl 4-hydroxylases that modify the hypoxia-inducible factor, *J. Biol. Chem.* 2003; 278:30772–30780
 65. **Ivan M, Haberberger T, Gervasi DC, Michelson KS, Günzler V, Kondo K, Yang H, Sorokina I, Conaway RC, Conaway JW, Kaelin WG** Biochemical purification and pharmacological inhibition of a mammalian prolyl hydroxylase acting on hypoxia-inducible factor, *Proc. Natl. Acad. Sci. U.S.A.* 2002; 99:13459–13464
 66. **Hewitson KS, McNeill LA, Riordan MV, Tian Y, Bullock AN, Welford RW, Elkins JM, Oldham NJ, Bhattacharya S, Gleadle JM, Ratcliffe PJ, Pugh CW, Schofield CJ** Hypoxia-inducible factor (HIF) asparagine hydroxylase is identical to factor inhibiting HIF (FIH) and is related to the cupin structural family, *J. Biol. Chem.* 2002; 277:26351–26355
 67. **Lando D, Peet DJ, Gorman JJ, Whelan DA, Whitelaw ML, Bruick RK** FIH-1 is an asparaginyl hydroxylase enzyme that regulates the transcriptional activity of hypoxia-inducible factor, *Genes Dev.* 2002; 16:1466–1471
 68. **Lando D, Peet DJ, Whelan DA, Gorman JJ, Whitelaw ML** Asparagine hydroxylation of the HIF transactivation domain a hypoxic switch, *Science.* 2002; 295:858–861
 69. **Manalo DJ, Rowan A, Lavoie T, Natarajan L, Kelly BD, Ye SQ, Garcia JGN, Semenza GL** Transcriptional regulation of vascular endothelial cell responses to hypoxia by HIF-1, *Blood.* 2005; 105:659–669
 70. **Wenger RH, Stiehl DP, Camenisch G** Integration of oxygen signaling at the consensus HRE, *Sci. STKE.* 2005; 2005:re12

71. **Gu YZ, Moran SM, Hogenesch JB, Wartman L, Bradfield CA** Molecular characterization and chromosomal localization of a third alpha-class hypoxia inducible factor subunit, HIF3alpha, *Gene Expr.* 1998; 7:205–213
72. **Hara S, Hamada J, Kobayashi C, Kondo Y, Imura N** Expression and characterization of hypoxia-inducible factor (HIF)-3alpha in human kidney: suppression of HIF-mediated gene expression by HIF-3alpha, *Biochem. Biophys. Res. Commun.* 2001; 287:808–813
73. **Maynard MA, Evans AJ, Hosomi T, Hara S, Jewett, Michael A S, Ohh M** Human HIF-3alpha4 is a dominant-negative regulator of HIF-1 and is down-regulated in renal cell carcinoma, *FASEB J.* 2005; 19:1396–1406
74. **Maynard MA, Qi H, Chung J, Lee, Eric H L, Kondo Y, Hara S, Conaway RC, Conaway JW, Ohh M** Multiple splice variants of the human HIF-3 alpha locus are targets of the von Hippel-Lindau E3 ubiquitin ligase complex, *J. Biol. Chem.* 2003; 278:11032–11040
75. **Prabhakar NR, Semenza GL** Adaptive and maladaptive cardiorespiratory responses to continuous and intermittent hypoxia mediated by hypoxia-inducible factors 1 and 2, *Physiol. Rev.* 2012; 92:967–1003
76. **Pasanen A, Heikkilä M, Rautavuoma K, Hirsilä M, Kivirikko KI, Myllyharju J** Hypoxia-inducible factor (HIF)-3alpha is subject to extensive alternative splicing in human tissues and cancer cells and is regulated by HIF-1 but not HIF-2, *Int. J. Biochem. Cell Biol.* 2010; 42:1189–1200
77. **Heikkilä M, Pasanen A, Kivirikko KI, Myllyharju J** Roles of the human hypoxia-inducible factor (HIF)-3 α variants in the hypoxia response, *Cell. Mol. Life Sci.* 2011; 68:3885–3901
78. **Zhang P, Yao Q, Lu L, Li Y, Chen P, Duan C** Hypoxia-inducible factor 3 is an oxygen-dependent transcription activator and regulates a distinct transcriptional response to hypoxia, *Cell Rep.* 2014; 6:1110–1121
79. **Hatanaka M, Shimba S, Sakaue M, Kondo Y, Kagechika H, Kokame K, Miyata T, Hara S** Hypoxia-inducible factor-3alpha functions as an accelerator of 3T3-L1 adipose differentiation, *Biol. Pharm. Bull.* 2009; 32:1166–1172
80. **Heidbreder M, Qadri F, Jöhren O, Dendorfer A, Depping R, Fröhlich F, Wagner KF, Dominiak P** Non-hypoxic induction of HIF-3alpha by 2-deoxy-D-glucose and insulin, *Biochem. Biophys. Res. Commun.* 2007; 352:437–443
81. **Goldberg AD, Allis CD, Bernstein E** Epigenetics: a landscape takes shape, *Cell.* 2007; 128:635–638
82. **Jaenisch R, Bird A** Epigenetic regulation of gene expression: how the genome integrates intrinsic and environmental signals, *Nat. Genet.* 2003; 33 Suppl:245–254
83. **Jenuwein T, Allis CD** Translating the histone code, *Science.* 2001; 293:1074–1080
84. **Kouzarides T** Chromatin modifications and their function, *Cell.* 2007; 128:693–705

85. **Jones PA, Takai D** The role of DNA methylation in mammalian epigenetics, *Science*. 2001; 293:1068–1070
86. **Mercer TR, Mattick JS** Structure and function of long noncoding RNAs in epigenetic regulation, *Nat. Struct. Mol. Biol.* 2013; 20:300–307
87. **Bird AP** Functions for DNA methylation in vertebrates, *Cold Spring Harb. Symp. Quant. Biol.* 1993; 58:281–285
88. **Bestor TH** The DNA methyltransferases of mammals, *Human Molecular Genetics*. 2000; 9:2395–2402
89. **Jin B, Li Y, Robertson KD** DNA methylation: superior or subordinate in the epigenetic hierarchy?, *Genes Cancer*. 2011; 2:607–617
90. **Meng H, Chen G, Gao H, Song X, Shi Y, Cao L** The emerging nexus of active DNA demethylation and mitochondrial oxidative metabolism in post-mitotic neurons, *Int J Mol Sci*. 2014; 15:22604–22625
91. **Jones PA** Functions of DNA methylation: islands, start sites, gene bodies and beyond, *Nat. Rev. Genet.* 2012; 13:484–492
92. **Goll MG, Bestor TH** Eukaryotic cytosine methyltransferases, *Annu. Rev. Biochem.* 2005; 74:481–514
93. **Laurent L, Wong E, Li G, Huynh T, Tsirigos A, Ong CT, Low HM, Kin Sung, Ken Wing, Rigoutsos I, Loring J, Wei C** Dynamic changes in the human methylome during differentiation, *Genome Res*. 2010; 20:320–331
94. **Lopez-Serra P, Esteller M** DNA methylation-associated silencing of tumor-suppressor microRNAs in cancer, *Oncogene*. 2012; 31:1609–1622
95. **Maunakea AK, Nagarajan RP, Bilenky M, Ballinger TJ, D'Souza C, Fouse SD, Johnson BE, Hong C, Nielsen C, Zhao Y, Turecki G, Delaney A, Varhol R, Thiessen N, Shchors K, Heine VM, Rowitch DH, Xing X, Fiore C, Schillebeeckx M, Jones, Steven J M, Haussler D, Marra MA, Hirst M, Wang T, Costello JF** Conserved role of intragenic DNA methylation in regulating alternative promoters, *Nature*. 2010; 466:253–257
96. **Drong AW, Lindgren CM, McCarthy MI** The genetic and epigenetic basis of type 2 diabetes and obesity, *Clin. Pharmacol. Ther.* 2012; 92:707–715
97. **Herrera BM, Keildson S, Lindgren CM** Genetics and epigenetics of obesity, *Maturitas*. 2011; 69:41–49
98. **Speliotes EK, Willer CJ, Berndt SI, Monda KL, Thorleifsson G, Jackson AU, Lango Allen H, Lindgren CM, Luan J, Mägi R, Randall JC, Vedantam S, Winkler TW, Qi L, Workalemahu T, Heid IM, Steinthorsdottir V, Stringham HM, Weedon MN, Wheeler E, Wood AR, Ferreira T, Weyant RJ, Segrè AV, Estrada K, Liang L, Nemesh J, Park J, Gustafsson S, Kilpeläinen TO, Yang J, Bouatia-Naji N, Esko T, Feitosa MF, Kutalik Z, Mangino M, Raychaudhuri S, Scherag A, Smith AV, Welch R, Zhao JH, Aben KK, Absher DM, Amin N, Dixon**

AL, Fisher E, Glazer NL, Goddard ME, Heard-Costa NL, Hoesel V, Hottenga J, Johansson A, Johnson T, Ketkar S, Lamina C, Li S, Moffatt MF, Myers RH, Narisu N, Perry JRB, Peters MJ, Preuss M, Ripatti S, Rivadeneira F, Sandholt C, Scott LJ, Timpson NJ, Tyrer JP, van Wingerden S, Watanabe RM, White CC, Wiklund F, Barlassina C, Chasman DI, Cooper MN, Jansson J, Lawrence RW, Pellikka N, Prokopenko I, Shi J, Thiering E, Alavere H, Alibrandi MTS, Almgren P, Arnold AM, Aspelund T, Atwood LD, Balkau B, Balmforth AJ, Bennett AJ, Ben-Shlomo Y, Bergman RN, Bergmann S, Biebermann H, Blakemore AIF, Boes T, Bonnycastle LL, Bornstein SR, Brown MJ, Buchanan TA, Busonero F, Campbell H, Cappuccio FP, Cavalcanti-Proença C, Chen YI, Chen C, Chines PS, Clarke R, Coin L, Connell J, Day INM, den Heijer M, Duan J, Ebrahim S, Elliott P, Elosua R, Eiriksdottir G, Erdos MR, Eriksson JG, Facheris MF, Felix SB, Fischer-Posovszky P, Folsom AR, Friedrich N, Freimer NB, Fu M, Gaget S, Gejman PV, Geus EJC, Gieger C, Gjesing AP, Goel A, Goyette P, Grallert H, Grässler J, Greenawalt DM, Groves CJ, Gudnason V, Guiducci C, Hartikainen A, Hassanali N, Hall AS, Havulinna AS, Hayward C, Heath AC, Hengstenberg C, Hicks AA, Hinney A, Hofman A, Homuth G, Hui J, Igl W, Iribarren C, Isomaa B, Jacobs KB, Jarick I, Jewell E, John U, Jørgensen T, Jousilahti P, Jula A, Kaakinen M, Kajantie E, Kaplan LM, Kathiresan S, Kettunen J, Kinnunen L, Knowles JW, Kolcic I, König IR, Koskinen S, Kovacs P, Kuusisto J, Kraft P, Kvaløy K, Laitinen J, Lantieri O, Lanzani C, Launer LJ, Lecoeur C, Lehtimäki T, Lettre G, Liu J, Lokki M, Lorentzon M, Luben RN, Ludwig B, Manunta P, Marek D, Marre M, Martin NG, McArdle WL, McCarthy A, McKnight B, Meitinger T, Melander O, Meyre D, Midthjell K, Montgomery GW, Morken MA, Morris AP, Mulic R, Ngwa JS, Nelis M, Neville MJ, Nyholt DR, O'Donnell CJ, O'Rahilly S, Ong KK, Oostra B, Paré G, Parker AN, Perola M, Pichler I, Pietiläinen KH, Platou CGP, Polasek O, Pouta A, Rafelt S, Raitakari O, Rayner NW, Ridderstråle M, Rief W, Ruukonen A, Robertson NR, Rzehak P, Salomaa V, Sanders AR, Sandhu MS, Sanna S, Saramies J, Savolainen MJ, Scherag S, Schipf S, Schreiber S, Schunkert H, Silander K, Sinisalo J, Siscovick DS, Smit JH, Soranzo N, Sovio U, Stephens J, Surakka I, Swift AJ, Tammesoo M, Tardif J, Teder-Laving M, Teslovich TM, Thompson JR, Thomson B, Tönjes A, Tuomi T, van Meurs, Joyce B J, van Ommen G, Vatin V, Viikari J, Visvikis-Siest S, Vitart V, Vogel CIG, Voight BF, Waite LL, Wallaschofski H, Walters GB, Widen E, Wiegand S, Wild SH, Willemsen G, Witte DR, Witteman JC, Xu J, Zhang Q, Zgaga L, Ziegler A, Zitting P, Beilby JP, Farooqi IS, Hebebrand J, Huikuri HV, James AL, Kähönen M, Levinson DF, Macciardi F, Nieminen MS, Ohlsson C, Palmer LJ, Ridker PM, Stumvoll M, Beckmann JS, Boeing H, Boerwinkle E, Boomsma DI, Caulfield MJ, Chanock

- SJ, Collins FS, Cupples LA, Smith GD, Erdmann J, Froguel P, Grönberg H, Gyllensten U, Hall P, Hansen T, Harris TB, Hattersley AT, Hayes RB, Heinrich J, Hu FB, Hveem K, Illig T, Jarvelin M, Kaprio J, Karpe F, Khaw K, Kiemeny LA, Krude H, Laakso M, Lawlor DA, Metspalu A, Munroe PB, Ouwehand WH, Pedersen O, Penninx BW, Peters A, Pramstaller PP, Quertermous T, Reinehr T, Rissanen A, Rudan I, Samani NJ, Schwarz PEH, Shuldiner AR, Spector TD, Tuomilehto J, Uda M, Uitterlinden A, Valle TT, Wabitsch M, Waeber G, Wareham NJ, Watkins H, Wilson JF, Wright AF, Zillikens MC, Chatterjee N, McCarroll SA, Purcell S, Schadt EE, Visscher PM, Assimes TL, Borecki IB, Deloukas P, Fox CS, Groop LC, Haritunians T, Hunter DJ, Kaplan RC, Mohlke KL, O'Connell JR, Peltonen L, Schlessinger D, Strachan DP, van Duijn CM, Wichmann H, Frayling TM, Thorsteinsdottir U, Abecasis GR, Barroso I, Boehnke M, Stefansson K, North KE, McCarthy MI, Hirschhorn JN, Ingelsson E, Loos RJJ** Association analyses of 249,796 individuals reveal 18 new loci associated with body mass index, *Nat. Genet.* 2010; 42:937–948
99. **Schleinitz D, Böttcher Y, Blüher M, Kovacs P** The genetics of fat distribution, *Diabetologia.* 2014; 57:1276–1286
 100. **Feil R, Fraga MF** Epigenetics and the environment: emerging patterns and implications, *Nat. Rev. Genet.* 2011; 13:97–109
 101. **Lienert F, Wirbelauer C, Som I, Dean A, Mohn F, Schübeler D** Identification of genetic elements that autonomously determine DNA methylation states, *Nat. Genet.* 2011; 43:1091–1097
 102. **Schadt EE** Molecular networks as sensors and drivers of common human diseases, *Nature.* 2009; 461:218–223
 103. **Li M, Wu H, Wang T, Xia Y, Jin L, Jiang A, Zhu L, Chen L, Li R, Li X** Co-methylated genes in different adipose depots of pig are associated with metabolic, inflammatory and immune processes, *Int. J. Biol. Sci.* 2012; 8:831–837
 104. **Demerath EW, Guan W, Grove ML, Aslibekyan S, Mendelson M, Zhou Y, Hedman ÅK, Sandling JK, Li L, Irvin MR, Zhi D, Deloukas P, Liang L, Liu C, Bressler J, Spector TD, North K, Li Y, Absher DM, Levy D, Arnett DK, Fornage M, Pankow JS, Boerwinkle E** Epigenome-wide association study (EWAS) of BMI, BMI change and waist circumference in African American adults identifies multiple replicated loci, *Human Molecular Genetics.* 2015
 105. **Dick KJ, Nelson CP, Tsaprouni L, Sandling JK, Aïssi D, Wahl S, Meduri E, Morange P, Gagnon F, Grallert H, Waldenberger M, Peters A, Erdmann J, Hengstenberg C, Cambien F, Goodall AH, Ouwehand WH, Schunkert H, Thompson JR, Spector TD, Gieger C, Trégouët D, Deloukas P, Samani NJ** DNA methylation and body-mass index: a genome-wide analysis, *Lancet.* 2014; 383:1990–1998

-
106. **Rönn T, Volkov P, Gillberg L, Kokosar M, Perfilyev A, Jacobsen AL, Jørgensen SW, Brøns C, Jansson P, Eriksson K, Pedersen O, Hansen T, Groop L, Stener-Victorin E, Vaag A, Nilsson E, Ling C** Impact of age, BMI and HbA1c levels on the genome-wide DNA methylation and mRNA expression patterns in human adipose tissue and identification of epigenetic biomarkers in blood, *Human Molecular Genetics*. 2015
 107. **Landegren U, Nilsson M, Kwok PY** Reading bits of genetic information: methods for single-nucleotide polymorphism analysis, *Genome Res*. 1998; 8:769–776
 108. **Cooper DN, Smith BA, Cooke HJ, Niemann S, Schmidtke J** An estimate of unique DNA sequence heterozygosity in the human genome, *Hum. Genet*. 1985; 69:201–205
 109. **Kwok PY, Deng Q, Zakeri H, Taylor SL, Nickerson DA** Increasing the information content of STS-based genome maps: identifying polymorphisms in mapped STSs, *Genomics*. 1996; 31:123–126
 110. **Graur D, Li W** Fundamentals of molecular evolution. 2nd edn. Sunderland, Mass.: Sinauer Associates; 2000
 111. **Bell JT, Pai AA, Pickrell JK, Gaffney DJ, Pique-Regi R, Degner JF, Gilad Y, Pritchard JK** DNA methylation patterns associate with genetic and gene expression variation in HapMap cell lines, *Genome Biol*. 2011; 12:R10
 112. **Boks MP, Derks EM, Weisenberger DJ, Strengman E, Janson E, Sommer IE, Kahn RS, Ophoff RA** The relationship of DNA methylation with age, gender and genotype in twins and healthy controls, *PLoS ONE*. 2009; 4:e6767
 113. **Swinburn BA, Sacks G, Hall KD, McPherson K, Finegood DT, Moodie ML, Gortmaker SL** The global obesity pandemic: shaped by global drivers and local environments, *Lancet*. 2011; 378:804–814
 114. **Trayhurn P, Alomar SY** Oxygen deprivation and the cellular response to hypoxia in adipocytes - perspectives on white and brown adipose tissues in obesity, *Front Endocrinol (Lausanne)*. 2015; 6:19